arXiv:math/9904099v1 [math.AG] 20 Apr 1999

Stokes Matrices and Monodromy of the Quantum Cohomology of Projective Spaces

Davide Guzzetti

International School for Advanced Studies (SISSA - ISAS)
Via Beirut, 2–4, 34014 Trieste, Italy.
E-mail: guzzetti@sissa.it, Fax: +39 040-3787-528

August 1998

Abstract

In this paper we compute Stokes matrices and monodromy for the quantum cohomology of projective spaces. This problem can be formulated in a "classical" framework, as the problem of computation of Stokes matrices and monodromy of (systems of) differential equations with regular and irregular singularities. We prove that the Stokes' matrix of the quantum cohomology coincides with the Gram matrix in the theory of derived categories of coherent sheaves. We also study the monodromy group of the quantum cohomology and we show that it is related to hyperbolic triangular groups.

1 Introduction

In this paper we compute Stokes' matrices and monodromy group for the Frobenius manifold given by the quantum cohomology of the projective space $\mathbf{CP}^d$. Our main motivation is to study the links between quantum cohomology and the theory of coherent sheaves.

Stokes matrices first appeared in the theory of WDVV equations of associativity in the paper [7] by B. Dubrovin. WDVV equations were formulated in a geometrical setting: the theory of Frobenius manifolds. From then on, the notion of Frobenius manifold has been largely studied in many papers, of which we cite [9], [10], [11].

The Stokes' matrix is a part of the *monodromy data* for a semisimple Frobenius manifold. Monodromy data can serve as natural moduli of semisimple Frobenius manifolds. More precisely, any local chart of the atlas covering the manifold is reconstructed from the monodromy data. The glueing of the local charts is described by the action of the braid group on the data, particularly, on the central connection matrix and on the Stokes' matrix [9] [10].

One well-known example of Frobenius manifolds is the quantum cohomology of smooth projective varieties [20] [19] [22] [21] .

It was conjectured [11] that the Stokes matrix for the quantum cohomology of a good Fano variety X is equal to the Gram matrix of the bilinear form $\chi(E,F):=\sum_k (-1)^k \dim Ext^k(E,F)$ computed on a full collection of exceptional objects in the derived category $Der^b(Coh(X))$ of coherent sheaves on X. More precisely, let $Der^b(Coh(X))$ be the derived category of coherent sheaves on a smooth projective variety X of dimension d. An object E of $Der^b(Coh(X))$ is called *exceptional* if $Ext^i(E,E)=0$ for $0<i<d$, $Ext^0(E,E)=\mathbf{C}$ and $Ext^d(E,E)$ is of the smallest dimension (if X is a projective space, then $Ext^d(E,E)=0$). A collection $\{E_1,...,E_s\}$ of exceptional objects is an *exceptional collection* if for any $1\le m<n\le s$ we have $Ext^i(E_n,E_m)=0$ for any $i\ge 0$, $Ext^i(E_m,E_n)=0$ for any $i\ge 0$ except possibly for one value of i. A *full exceptional collection* is an exceptional collection which generates $Der^b(Coh(X))$ as a triangulated category. This theory is developed in [23] [24] [5]. We say that a Fano variety is *good* if it has a full exceptional collection.

It is known that $X=\mathbf{CP}^d$ is good, the collection of sheaves on $\mathbf{CP}^d$ $\{\mathcal{O}(n)\}_{n\in\mathbf{Z}}$ is exceptional, and $\{E_1,E_2,...,E_{d+1}\}:=\{\mathcal{O},\mathcal{O}(1),\ ...,\mathcal{O}(d)\}$ is a full exceptional collection [3], [16]. In this case, $s_{ij}=\chi\left(\mathcal{O}(i-1),\mathcal{O}(j-1)\right)$, $i,j=1,2,...,d+1$ has the "canonical form":

$$s_{ij}=\begin{pmatrix} d+j-i \\ j-i \end{pmatrix} \qquad i<j$$

$$s_{ii}=1, \qquad s_{ij}=0 \quad i>j$$

The inverse to this matrix has entries a_{ij}

$$a_{ij} = (-1)^{j-i} \begin{pmatrix} d+1 \\ j-i \end{pmatrix} \qquad i<j$$

$$a_{ii}=1, \qquad a_{ij}=0 \quad i>j$$

This matrix is equivalent to the one above with respect to the action of the braid group. We will also call it "canonical".

The mentioned conjecture claims that the Stokes matrix of the quantum cohomology of $\mathbf{CP}^d$ is equal to the above Gram matrix (modulo the action of the braid group: remarkably, this action on the Stokes matrix for the Frobenius manifold coincides with the natural action of the braid group on the collections of exceptional objects [25] [23]).

This conjecture has its origin in the paper by Cecotti and Vafa [6], where another Stokes matrix introduced in [8] for the tt^* equations was found in the case of the $\mathbf{CP}^2$ topological σ model. It was suggested, on physical arguments, that the entries of the Stokes' matrix $S = \begin{pmatrix} 1 & x & y \\ 0 & 1 & z \\ 0 & 0 & 1 \end{pmatrix}$ are integers. They must satisfy a Diophantine equation $x^2+y^2+z^2-xyz=0$ whose integer solutions (x,y,z) are all equivalent to (3,3,3) modulo the action of the braid group. The authors of [6] also suggested that their matrix must coincide with the Stokes matrix defined in the theory of WDVV equations of associativity, that is, in the geometrical theory of Frobenius manifolds for 2D topological field theories [7], [9].

Later, in [25], the links between $N=2$ supersymmetric field theories and the theory of derived categories were further investigated and the coincidence of $\chi(E_i,E_j)$ with the Stokes matrix of tt^* for $\mathbf{CP}^d$ was conjectured.

The conjecture may probably be derived from more general conjectures by Kontsevich in the framework of categorical mirror symmetry. To my knowledge, the subject was discussed in [18] (I thank B. Dubrovin for this reference).

The main result of this paper is the proof (Theorem 2, 2′) that the conjecture about coincidence of the Stokes matrix for quantum cohomology of $\mathbf{CP}^d$ and the Gram matrix $\chi(E_i,E_j)$ of a full exceptional collection in $Der^b(Coh(\mathbf{CP}^d))$ is true. In this way, we generalize to any d the result obtained in [10] for $d=2$.

We remark that it has not yet been proved that the Stokes' matrix for tt^* equations and the Stokes' matrix for the corresponding Frobenius manifold coincide. This point deserves further investigation.

We also study the structure of the monodromy group of the quantum cohomology of $\mathbf{CP}^d$. The notion of *monodromy group* of a Frobenius manifold was introduced in [9]. We prove (Theorem 3) that for $d=3$ the group is isomorphic to the subgroup of orientation preserving transformations in the hyperbolic triangular group $[2,4,\infty]$. In [10] it was proved that for $d=2$ the monodromy group is isomorphic to the direct product of the subgroup of orientation preserving transformations in $[2,3,\infty]$ and the cyclic group of order 2, $C_2=\{\pm\}$. Our numerical calculations also suggest that for any d even the monodromy group may be isomorphic to the orientation preserving transformations in $[2,d+1,\infty]$, and for any d odd to the direct product of the orientation preserving transformations in $[2,d+1,\infty]$ by C_2.

Acknowledgments

I am indebted to Prof. B Dubrovin who introduced me to the problem and constantly gave me suggestions and advice. I also thank M. Bertola and M. Mazzocco for useful discussions.

2 The system corresponding to $\mathbf{CP}^{k-1}$

We introduce here the linear system of differential equations whose Stokes matrices are the Stokes matrices for the quantum cohomology of $\mathbf{CP}^{k-1}$ (we use the more convenient choice $k=d+1$).

In the quantum cohomology of $\mathbf{CP}^{k-1}$ we choose flat coordinate $t^1,..,t^k$ for the symmetric non degenerate *bilinear form* $<\ ,\ >$:

$$\eta_{\alpha\beta} = <\partial_\alpha, \partial_\beta> = \delta_{\alpha+\beta,k+1} \qquad \text{where } \partial_\alpha = \frac{\partial}{\partial t^\alpha}$$

Let η be the matrix $(\eta_{\alpha\beta})$. In flat coordinates the *Euler vector field* is

$$E = \sum_{\alpha\neq 2} (1-q_\alpha)t^\alpha \frac{\partial}{\partial t^\alpha} + \ k\frac{\partial}{\partial t^2}$$

$$q_1 = 0,\ q_2 = 1,\ q_3 = 2,\ ...,\ q_k = k-1$$

the *multiplication* is

$$\partial_\alpha \cdot \partial_\beta = c_{\alpha\beta}^{\gamma}(t)\partial_\gamma \qquad \text{where} \quad c_{\alpha\beta\gamma}(t) := \eta_{\delta\gamma}c^{\delta}_{\alpha\beta}(t) = \partial_\alpha\partial_\beta\partial_\gamma F(t)$$

$$F(t) = \frac{1}{6}\sum_{\alpha+\beta+\gamma=k+2} t^\alpha t^\beta t^\gamma + \sum_{l=1}^{\infty}\Phi_l(t^3,...,t^k)e^{lt^2}$$

$$\Phi_l(t^3,...,t^k) = \sum_{n=2}^{\infty}\left[\sum_{\alpha_1+...+\alpha_n=(l+1)(k-1)+l-1+n} I(l;\alpha_1,...,\alpha_n)\frac{t^{\alpha_1}...t^{\alpha_n}}{n!}\right], \qquad I(1,k,k)=1$$

and the *unity vector field* is $e = \frac{\partial}{\partial t^1}$. Finally, let

$$\mu = \text{diag}(\mu_1,...,\mu_k) = \text{diag}(-\frac{k-1}{2},-\frac{k-3}{2},...,\frac{k-3}{2},\frac{k-1}{2}), \qquad \mu_\alpha = q_\alpha - \frac{d}{2}$$

Consider the system of differential equations determining deformed flat coordinates (see [9] [10]):

$$\partial_z\xi = (\mathcal{U}(t)+\frac{1}{z}\mu)\xi \tag{1}$$

$$\partial_\alpha\xi = zC_\alpha(t)\xi$$

where $z\in \mathbf{C}$, $\partial_z := \frac{\partial}{\partial z}$ and ξ is a column vector of components

$$\xi^\alpha = \eta^{\alpha\beta}\partial_\beta\tilde{t}(t,z)$$

Here $(\eta^{\alpha\beta}) = (\eta_{\mu\nu})^{-1}$ and $\tilde{t}(t,z)$ is one of the k (deformed) flat coordinates. $\mathcal{U}(t)$ is the matrix of multiplication by the Euler vector field $E(t)$, and $C_\alpha(t)$ is the matrix of entries $(C_\alpha)^{\beta}_{\ \gamma} := c^{\beta}_{\alpha\gamma}$. The monodromy data of the system (1) are, by definition, the *monodromy data* of the quantum cohomology of $\mathbf{CP}^{k-1}$ in the local chart containing t. Let us compute $C_2(t)$ and $\mathcal{U}(t)$ at the semisimple point $(0,t^2,0,...,0)$:

$$E\cdot\ \partial_\beta = E^\gamma c^{\alpha}_{\gamma\beta}\ \partial_\alpha = E^2 c^{\alpha}_{2\beta}\ \partial_\alpha =$$
$$= k\ c^{\alpha}_{2\beta}\partial_\alpha \equiv \mathcal{U}^{\alpha}_{\ \beta}\partial_\alpha$$

Moreover $c_{2\alpha\beta}(0,t^2,0,...,0) = \partial_2\partial_\alpha\partial_\beta F(0,t^2,...,0)$. This immediately yields

$$C_2(0,t^2,0,...,0) = \begin{pmatrix} 0 & & & & e^{t^2} \\ 1 & 0 & & & \\ & 1 & 0 & & \\ & & \ddots & \ddots & \\ & & & 1 & 0 \end{pmatrix}, \quad \mathcal{U}(0,t^2,0,...,0) = \begin{pmatrix} 0 & & & & ke^{t^2} \\ k & 0 & & & \\ & k & 0 & & \\ & & \ddots & \ddots & \\ & & & k & 0 \end{pmatrix}$$

Let $y_\alpha := \eta_{\alpha\beta}\xi^\beta \equiv \partial_\alpha\tilde{t}$. It satisfies

$$\partial_z y = \left(\hat{\mathcal{U}}(t^2) - \frac{1}{z}\mu\right)y \tag{2}$$

$$\partial_2 y = z\hat{C}_2(t^2)y \tag{3}$$

where

$$\hat{C}_2(t^2) := \eta\ C_2(0,t^2,...,0)\ \eta^{-1} = \begin{pmatrix} 0 & 1 & & & & \\ & 0 & 1 & & & \\ & & 0 & 1 & & \\ & & & \ddots & \ddots & \\ & & & & 0 & 1 \\ e^{t^2} & & & & & 0 \end{pmatrix}$$

$$\hat{\mathcal{U}}(t^2) := \eta\, \mathcal{U}(0, t^2, ..., 0)\ \eta^{-1} = \begin{pmatrix} 0 & k & & & & \\ & 0 & k & & & \\ & & 0 & k & & \\ & & & \ddots & \ddots & \\ & & & & 0 & k \\ ke^{t^2} & & & & & 0 \end{pmatrix}$$

Lemma 1: *Let $y(t^2, z) = (y_1(t^2, z), ..., y_k(t^2, z))^T$ be a column vector solution of the above system (2) (3). With the following substitution*

$$y_\alpha(t^2, z) = \frac{1}{k^{\alpha-1}}\ z^{\frac{k-1}{2}-\alpha+1}\ (z\partial_z)^{(\alpha-1)}\ \varphi(t^2, z) \tag{4}$$

$$\equiv z^{\frac{k-1}{2}-\alpha+1}\ \partial_2^{\alpha-1}\varphi(t^2, z), \quad \alpha = 1, 2, ..., k \tag{5}$$

the above system is equivalent to the equations

$$(z\partial_z)^k\varphi = (kz)^k e^{t_2}\ \varphi \tag{6}$$

$$\partial_2^k\varphi = z^k e^{t_2}\ \varphi \tag{7}$$

The proof is a simple calculation we leave to the reader.

The substitution of the lemma implies $\partial_2\varphi = \frac{1}{k}z\partial_z\varphi$. Then

$$e^{\frac{t^2}{k}}\frac{\partial}{\partial e^{\frac{t^2}{k}}}\varphi(t^2, z) = z\frac{\partial}{\partial z}\varphi(t^2, z)$$

which implies (with abuse of notation)

$$\varphi(t^2, z) \equiv \varphi(ze^{\frac{t^2}{k}})$$

Namely, φ (at $(0, t^2, ..., 0)$) depends on one argument $w = ze^{\frac{t^2}{k}}$ and satisfies the *generalized hypergeometric equation*

$$\left(w\frac{d}{dw}\right)^k \varphi(w) = (kw)^k\ \varphi(w) \tag{8}$$

The equation is equivalent to the system

$$\frac{dY}{dw} = \left[\hat{\mathcal{U}} + \frac{\hat{\mu}}{w}\right] Y \tag{9}$$

where

$$Y_n(w) = \frac{1}{k^{n-1}}\ w^{\frac{k-1}{2}-n+1}\ (w\partial_w)^{(n-1)}\ \varphi(w) \quad n = 1, 2, ..., k \tag{10}$$

$$\hat{\mathcal{U}} := \hat{\mathcal{U}}(0) = \begin{bmatrix} 0 & k & & & & \\ & 0 & k & & & \\ & & 0 & k & & \\ & & & \ddots & \ddots & \\ & & & & \ddots & k \\ k & & & & & 0 \end{bmatrix}$$

$$\hat{\mu} := -\mu = \ \mathrm{diag}\left(\frac{k-1}{2}, \frac{k-3}{2}, \frac{k-5}{2}, ..., -\frac{k-3}{2}, -\frac{k-1}{2}\right)$$

The system (9) may also be interpreted as the system (2) with $t^2 = 0$. We will return later (section 4) on the connection between its monodromy data and the monodromy data of the system (2).

Let us study system (9). We change notation and choose the more familiar letter z instead of w. So, the system (9) is re-written as

$$\frac{dY}{dz} = \left[\hat{\mathcal{U}} + \frac{\hat{\mu}}{z}\right] Y \tag{11}$$

and (10) (8) become

$$Y_n(z) = \frac{1}{k^{n-1}}\ z^{\frac{k-1}{2}-n+1}\ (z\partial_z)^{(n-1)}\ \varphi(z) \quad n = 1, 2, ..., k \tag{12}$$

$$\left(z\frac{d}{dz}\right)^k \varphi(z) = (kz)^k\ \varphi(z) \tag{13}$$

The point $z = 0$ is e fuchsian singularity, and $z = \infty$ is a singularity of the second kind. (11) has a fundamental matrix solution $Y_0(z)$ whose behaviour at $z = 0$ is

$$Y_0(z) = (I + O(z)) z^{\hat{\mu}} z^R \qquad R = \begin{bmatrix} 0 & k & & & & \\ & 0 & k & & & \\ & & 0 & k & & \\ & & & \ddots & \ddots & \\ & & & & \ddots & k \\ & & & & & 0 \end{bmatrix}$$

and the monodromy for a counter-clockwise loop around the origin is $e^{2\pi i(\hat{\mu}+R)}$.

The characteristic polynomial of the matrix $\hat{\mathcal{U}}$ is $0 = \det(\hat{\mathcal{U}} - u) = (-u)^k + (-1)^{k+1}k^k$. It has k distinct eigenvalues $u_n = k\ e^{\frac{2\pi i(n-1)}{k}}$, $n = 1, ..., k$. The equations for the eigenvector $\mathbf{x}_n$ corresonding to u_n, namely $\hat{\mathcal{U}}\mathbf{x}_n = u_n\ \mathbf{x}_n$, written for the components ${x^1}_n$, ...,${x^k}_n$ of the column vector $\mathbf{x}_n$ are

$${x^{l+1}}_n = e^{\frac{2\pi i(n-1)}{k}}\ {x^l}_n \qquad l = 1, 2, ..., k-1$$

$${x^1}_n = e^{\frac{2\pi i(n-1)}{k}} {x^k}_n$$

With the choice ${x^1}_n = e^{\frac{i\pi(n-1)}{k}}$ we get $\mathbf{x}_n = (e^{\frac{i\pi(n-1)}{k}}, e^{\frac{i3\pi(n-1)}{k}}, e^{\frac{i5\pi(n-1)}{k}}, ..., e^{-\frac{i\pi(n-1)}{k}})^T$ (T stands for transpose). The matrix

$$X = \frac{1}{\sqrt{k}}\ [\mathbf{x}_1|\mathbf{x}_2|...|\mathbf{x}_k] = \frac{1}{\sqrt{k}}\ ({x^j}_n) \qquad {x^j}_n = e^{(2j-1)i\pi\frac{n-1}{k}} \quad j, n = 1, 2, ..., k$$

puts $\mathcal{U}$ in diagonal form:

$$U = X^{-1}\hat{\mathcal{U}}X = \mathrm{diag}(u_1, u_2, ..., u_n, ..., u_k) \quad u_n = k\ e^{\frac{2\pi i(n-1)}{k}}$$

We stress that $u_i \neq u_j$ for $i \neq j$. The system (11) is transformed by the gauge X in an equivalent form

$$\frac{d\tilde{Y}}{dz} = \left[U + \frac{V}{z}\right]\tilde{Y} \tag{14}$$

$$\tilde{Y} = X^{-1}Y, \quad U = X^{-1}\hat{\mathcal{U}}X, \quad V = X^{-1}\hat{\mu}X$$

Observe that

$$\eta\hat{\mu} + \hat{\mu}\eta = 0 \qquad XX^T = \eta^{-1}$$

This implies that V is skew-symmetric

$$V + V^T = 0$$

With the gauge X, $Y_0(z)$ transforms into

$$\tilde{Y}_0(z) = (X^{-1} + O(z))\ z^{\hat{\mu}}z^R, \quad z \to 0$$

3 Asymptotic Behaviour and Stokes' Phenomenon

Our aim is to explicitely compute a Stokes' matrix for the above system (14), or for the system (11). The system (14) has formal solution

$$\tilde{Y}_F = \left[I + \frac{F_1}{z} + \frac{F_2}{z^2} + ... \right] \; e^{z\;U}$$

where F_j's are $k \times k$ matrices. It is a well known result that fundamental matrix solutions exist which have $\tilde{Y}_F$ as asymptotic expansion for $z \to \infty$ in some "admissible" sectors of the complex plane of angular width greater then π. In order to find such sectors we need the so called *Stokes' rays*, defined by

$$\Re\; ((u_r - u_s)z) = 0, \quad r \neq s \quad \Rightarrow \quad \arg z = -\arg(u_r - u_s) + \frac{\pi}{2} + m\pi, \quad m = 0, 1$$

There exists a *unique* solution of the system asymptotic to $\tilde{Y}_F$ in a sector greater then π and bounded by the first two Stokes' rays we meet extending over π the angular width of the sector. The general theory of Stokes' phenomenon is found in the classical paper by W. Balser, W.B. Jurkat, D.A. Lutz [1]. Stokes matrices are also a natural monodromy datum in the theory of isomonodromy deformation [17] [13] [14] [15] [9] [12].

A possible choice for the labelling of the rays is the following: we call R_{rs} the Stokes' ray

$$R_{rs} = \{ z = -i\rho(\bar{u_r} - \bar{u_s}), \quad \rho > 0 \} \qquad r \neq s$$

Lemma 2: *For $r < s$ the Stokes' rays of the system (14) are*

$$R_{rs} = \left\{ z = \rho\;\; \exp\left(i\left[\frac{2\pi}{k} - \frac{\pi}{k}(r+s) \right] \right), \quad \rho > 0 \right\}$$

$$R_{sr} = -R_{rs}$$

Proof: Just compute

$$-i(\bar{u_r} - \bar{u_s}) = -i(e^{-i\frac{2\pi}{k}(r-1)} - e^{-i\frac{2\pi}{k}(s-1)}) =$$
$$= 2\sin\left(\frac{\pi}{k}(s-r)\right)\;\; e^{\left(i\left[\frac{2\pi}{k} - \frac{\pi}{k}(r+s)\right]\right)}$$

Then we note that $\sin\left(\frac{\pi}{k}(s-r)\right)$ is positive because $0 < s - r \leq k - 1$.

□

Remark 1: $R_{rs} = R_{pq}$ for $r + s = p + q$. R_{12} is at $\arg z = -\frac{\pi}{k}$, R_{13} is at $\arg z = -\frac{2\pi}{k}$, and so on. For $r + s = k + 2$ the corresponding R_{rs}'s are at $\arg z = -\pi$ and the R_{sr}'s are at $\arg z = 0$. $R_{k-1,k}$ is at the angle $-2\pi + \frac{3\pi}{k}$ or, equivalently, at $\frac{3\pi}{k}$. See figure 1.

We choose two "admissible" overlapping sectors in a canonical way. Let l be an "admissible" line through the origin, namely a line not containing Stokes' rays. For our purposes we take

$$l = \{ z \mid z = \rho e^{i\epsilon}, \quad \rho \in \mathbf{R}, \quad 0 < \epsilon < \frac{\pi}{k} \}$$

l has a natural orientation inherited from $\mathbf{R}$. We call Π_R and Π_L the half planes to the right/left of l w.r.t its orientation.

$$\Pi_R = \{ -\pi + \epsilon < \arg z < \epsilon \} \quad \Pi_L = \{ \epsilon < \arg z < \pi + \epsilon \}$$

We then define two different "admissible" sectors $\mathcal{S}_L$, $\mathcal{S}_R$ which contain l

$$\mathcal{S}_L = \{ z \in \mathbf{C} \mid 0 < argz < \pi + \frac{\pi}{k} \} \supset \Pi_L$$

$$\mathcal{S}_R = \{ z \in \mathbf{C} \mid -\pi < argz < \frac{\pi}{k} \} \supset \Pi_R$$



k even k odd

Figure 1: Stokes' rays

We call che corresponding solutions $\tilde{Y}_L(z)$ and $\tilde{Y}_R(z)$.

Definition: The *Stokes' matrix* of the system (14) with respect to the admissible line l is the connection matrix S such that

$$\tilde{Y}_L(z) = \tilde{Y}_R(z) S \qquad 0 < \arg z < \frac{\pi}{k}$$

On the opposite overlapping region one can prove (as a consequence of the skew-symmetry of V, see [9]) that

$$\tilde{Y}_L(z) = \tilde{Y}_R(z\ e^{-2\pi i}) S^T \qquad \pi < \arg z < \pi + \frac{\pi}{k}$$

In [1] S is called a " Stokes' multiplier". The terminology in this field changes from one author to the other...

Definition: We call *central connection matrix* the connection matrix C such that

$$\tilde{Y}_0(z) = \tilde{Y}_R(z) C \quad z \in \Pi_R$$

It is clear that the system (11) has solutions $Y_0(z) = X\tilde{Y}_0(z)$, and $Y_L(z) = X\tilde{Y}_L(z)$, $Y_R(z) = X\tilde{Y}_R(z)$ asymptotic to $X\tilde{Y}_F(z)$ as $z \to \infty$ in $\mathcal{S}_L$ and $\mathcal{S}_R$ respectively, which are connected by the same S and C.

In order to compute the entries of S explicitely, we use the reduction of (11) to the generalized hypergeometric equation (13). If $\varphi^{(1)}(z)$, ..., $\varphi^{(k)}(z)$ is a basis of k linearly independent solutions of (13), then the matrix $Y(z)$ of entries (n,j) defined by

$$Y_n^{(j)}(z) := \frac{1}{k^{n-1}}\ z^{\frac{k-1}{2}-n+1}\ (z\partial_z)^{(n-1)}\ \varphi^{(j)}(z) \tag{15}$$

is a fundamental matrix for (11).

Lemma 3: *The generalized hypergeometric equation (13) has two bases of linerly independent solutions* $\varphi_{L/R}^{(1)}(z)$, *...,* $\varphi_{L/R}^{(k)}(z)$ *having asymptotic behaviours*

$$\varphi_{L/R}^{(n)} = \frac{1}{\sqrt{k}} \frac{e^{i\frac{\pi}{k}(n-1)}}{z^{\frac{k-1}{2}}}\ \exp\left[k\ e^{i\frac{2\pi}{k}(n-1)}\ z\right]\ \left(1 + O\left(\frac{1}{z}\right)\right), \qquad z \to \infty$$

in $\mathcal{S}_L$ *and* $\mathcal{S}_R$ *respectively. Let* $\Phi(z)$ *denote the row vector* $[\varphi^{(1)}(z), ..., \varphi^{(k)}(z)]$*. The fundamental matrices* $Y_L(z)$*,* $Y_R(z)$ *of (11) are expressed through formula (15) in terms of* $\Phi_L(z)$ *and* $\Phi_R(z)$ *and*

$$Y_L(z) = Y_R(z)\ S \qquad 0 < \arg z < \frac{\pi}{k}$$

if and only if

$$\Phi_L(z) = \Phi_R(z) S \qquad 0 < \arg z < \frac{\pi}{k}$$

Proof: Simply observe that for a fundamental solution in $\mathcal{S}_L$ or $\mathcal{S}_R$ (we omit subscripts L, R)

$$Y(z) = \begin{bmatrix} z^{\frac{k-1}{2}}\varphi^{(1)} & ... & z^{\frac{k-1}{2}}\varphi^{(k)} \\ \vdots & ... & \vdots \end{bmatrix} = X\tilde{Y}$$

which is asymptotic, for $z \to \infty$, to

$$\sim \begin{bmatrix} 1 & e^{i\frac{\pi}{k}} & e^{i\frac{2\pi}{k}} & e^{i\frac{3\pi}{k}} & ... & e^{i\frac{k-1}{k}\pi} \\ \vdots & \vdots & \vdots & \vdots & ... & \vdots \end{bmatrix} \begin{bmatrix} \exp(kz) & & & \\ & \exp(ke^{\frac{2\pi i}{k}}z) & & \\ & & \ddots & \\ & & & \exp(ke^{\frac{2\pi i(k-1)}{k}}) \end{bmatrix}$$

Now , the first row of $Y(z)$ is $z^{\frac{k-1}{2}}\Phi(z)$

$\square$

The Stokes' matrix S has entries

$$s_{ii} = 1$$

$$s_{ij} = 0 \quad \text{if } R_{ij} \subset \Pi_R$$

This follows from the fact that on the overlapping reagion $0 < \arg z < \frac{\pi}{k}$ there are no Stokes' rays and

$$e^{zU}\ S\ e^{-zU} \sim I, \qquad z \to \infty, \qquad \text{then} \quad e^{z(u_i - u_j)}\ s_{ij} \to \delta_{ij}$$

Moreover, $\Re\left(z(u_i - u_j)\right) > 0$ to the left of the ray R_{ij}, while $\Re\left(z(u_i - u_j)\right) < 0$ to the right (the natural orientation on R_{ij}, from $z = 0$ to ∞ is understood). This implies

$$|e^{zu_i}| > |e^{zu_j}| \qquad \text{and } e^{z(u_i - u_j)} \to \infty \quad \text{as } z \to \infty$$

on the left, while on the right

$$|e^{zu_i}| < |e^{zu_j}| \qquad \text{and } e^{z(u_i - u_j)} \to 0 \quad \text{as } z \to \infty$$

With this observations in mind, we prove the following

Lemma 4 : *S has a column whose entries are all zero but one. More precisely:*
For k even

$$s_{i,\frac{k}{2}+1} = 0 \quad \forall i \neq \frac{k}{2} + 1, \qquad s_{\frac{k}{2}+1,\frac{k}{2}+1} = 1$$

For k odd

$$s_{i,\frac{k+1}{2}} = 0 \quad \forall i \neq \frac{k+1}{2}, \qquad s_{\frac{k+1}{2},\frac{k+1}{2}} = 1$$

Proof: Let us determine n such that $s_{in} = 0$ for any $i \neq n$ and $s_{nn} = 1$. We need to find all rays in Π_R. We start with R_{rs} with $r < s$. We know that for $r + s = k + 2$ the ray is the negative real half-line (at angle $-\pi$). Then $R_{rs} \subset \Pi_R$ for $r + s \leq k + 1$ ($r < s$). Then, in Π_R we have

$$\begin{matrix} R_{12} & R_{13} & ... & R_{1k} \\ R_{23} & R_{24} & ... & R_{2,k-1} \\ R_{34} & R_{35} & ... & R_{3,k-2} \\ \vdots & \vdots & & \\ R_{ab} & & & \end{matrix}$$

where $R_{ab} = R_{\frac{k}{2},\frac{k}{2}+1}$ for k even, and $R_{\frac{k-1}{2},\frac{k+3}{2}}$ for k odd. In Π_R we have also R_{rs} with $r + s \geq k + 2$ and $r > s$. For fixed n we require $R_{in} \subset \Pi_R$ for any i. Namely,

$$\forall i < n \quad i + n \leq k + 1\ , \qquad \forall i > n \quad i + n \geq k + 2$$

This yields $n=\frac{k}{2}+1$ for k even, $n=\frac{k+1}{2}$ for k odd.

$\square$

Let $n(k)$ be $\frac{k}{2}+1$, or $\frac{k+1}{2}$. Lemma 4 implies that the $n(k)^{th}$ columns of Y_L and Y_R coincide. In particular, their asymptotic representation holds for $-\pi<\arg z<\pi+\frac{\pi}{k}$. Actually, this domain can be further enlarged, up to

$$-\frac{\pi}{k}-\pi<\arg z<\pi+\frac{\pi}{k} \qquad k \text{ even}$$
$$-\pi<\arg z<\pi+\frac{2\pi}{k} \qquad k \text{ odd}$$

To see this recall that $|e^{zu_i}|<|e^{zu_j}|$ on the right of R_{ij}, and conversely on the left. Then it is easy to see that for k even $\left|\exp(z\ u_{\frac{k}{2}+1})\right|$ dominates all exponentials in the sector $-\frac{\pi}{k}-\pi<\arg z<\frac{\pi}{k}-\pi$, while for k odd $\left|\exp(z\ u_{\frac{k+1}{2}})\right|$ dominates all exponentials in the sector $\pi<\arg z<\pi+\frac{2\pi}{k}$.

The first entry of the $n(k)^{th}$ column is $\varphi_L^{(n(k))}(z)\equiv\varphi_R^{(n(k))}(z)$ times $z^{\frac{k-1}{2}}$. Then $\varphi^{(n(k))}$ has the estabilished asymtotic behaviour on the enlarged domains above.

We now introduce an integral representation for a solution $\varphi(z)$ of the generalized hypergeometric equation which will allow us to compute the entries of S.

Lemma 5: *The function*

$$g^{(n)}(z)=\frac{1}{(2\pi)^{\frac{k+1}{2}}\ e^{i\pi(\frac{k}{2}-n-1)}}\int_{-c-i\infty}^{-c+i\infty} ds\ \Gamma^k(-s)\ e^{-i\pi k s}\ e^{i2(n-1)\pi s}\ z^{ks}$$

defined for $\frac{\pi}{2}-2(n-1)\frac{\pi}{k}<\arg z<\frac{3\pi}{2}-2(n-1)\frac{\pi}{k}$, $z\neq 0$ and for any positive number $c>0$, is a solution of the generalized hypergeometric equation (13) (the path of integration is a vertical line through $-c$). It has asymptotic behaviour

$$g^{(n)}(z)\sim\frac{1}{\sqrt{k}}\ \frac{e^{i\frac{\pi}{k}(n-1)}}{z^{\frac{k-1}{2}}}\ \exp\left(ke^{i\frac{2\pi}{k}(n-1)}z\right) \qquad z\to\infty$$

In particular, for $n(k)=\frac{k}{2}+1$ (k even), or $n(k)=\frac{k+1}{2}$ (k odd), the analytic continuation of $g^{(n(k))}(z)$ has the above asymptotic behaviour in the domains

$$-\pi-\frac{\pi}{k}<\arg z<\pi+\frac{\pi}{k} \qquad k \text{ even}$$
$$-\pi<\arg z<\pi+\frac{2\pi}{k} \qquad k \text{ odd}$$

and coincides with the solution $\varphi_L^{(n(k))}\equiv\varphi_R^{(n(k))}$ appearing in the first rows of the fundamental matrices Y_L and Y_R of the system (11).

The following identity holds

$$\sum_{m=0}^{k}\ (-1)^{m-k}\ \binom{k}{m}\ g^{(n)}(z\ e^{i\frac{2\pi}{k}m})=0 \tag{16}$$

We'll sketch the proof in Appendix 1.

Remark 2: Observe that for basic solutions of the hypergeometric equation $\Phi_{L/R}=[\varphi_{L/R}^{(1)},...,\varphi_{L/R}^{(k)}]$,

$$\varphi^{(n)}(z)\sim\ \frac{1}{\sqrt{k}}\frac{e^{i\frac{\pi}{k}(n-1)}}{z^{\frac{k-1}{2}}}\exp(ke^{i\frac{2\pi}{k}(n-1)}z)$$

on some sector, and

$$\varphi^{(n)}(ze^{\frac{2\pi i}{k}})\sim(-1)\ \frac{1}{\sqrt{k}}\frac{e^{i\frac{\pi}{k}([n+1]-1)}}{z^{\frac{k-1}{2}}}\exp(ke^{i\frac{2\pi}{k}([n+1]-1)}z)$$

like $-\varphi^{(n+1)}$, on the sector rotated by $\frac{-2\pi}{k}$. Note however that $\varphi^{(k)}(ze^{\frac{2\pi i}{k}})\sim(\sqrt{k}z^{\frac{k-1}{2}})^{-1}\ e^{kz}$, like $\varphi^{(1)}(z)$.

Also, note that $g^{(n)}(ze^{\frac{2\pi i}{k}})=-g^{(n+1)}(z)$ (we mean analytic continuations), with asymptotic behaviour on rotated domain.

4 Monodromy Data of the Quantum Cohomology of $\mathbf{CP}^{k-1}$

Let us return to the system (2).

$$\partial_z y = \left(\hat{\mathcal{U}}(t^2) + \frac{1}{z}\hat{\mu}\right) y \tag{17}$$

In this section we use the original notation $w = ze^{\frac{t^2}{k}}$. The system has a fundamental matrix

$$y_0(t^2,z) = (I + A_1(t^2)z + A_2(t^2)z^2 + ...)\ z^{\hat{\mu}}\ z^R, \qquad z \to 0$$

where R is the same of system (11). The series appearing in the solutions converges near $z = 0$. The matrix $\hat{\mathcal{U}}(t^2)$ has eigenvalues and eigenvectors

$$u_n(t^2) = e^{i\frac{2\pi}{k}(n-1)}\ e^{\frac{t^2}{k}} \ \equiv\ u_n e^{\frac{t^2}{k}} \qquad n = 1, ..., k$$

$$\mathbf{x}_n(t^2) \text{ of entries } {x^j}_n(t^2) = e^{i(2j-1)(n-1)\frac{\pi}{k}}\ e^{\frac{2j-1-k}{2k}t^2} \ \equiv {x^j}_n e^{\frac{2j-1-k}{2k}t^2}$$

Let $X(t^2) = ({x^j}_n(t^2))$. With the gauge $y = X(t^2)\ \tilde{y}(t^2,z)$ we obtain the equivalent system

$$\partial_z \tilde{y} = \left[U(t^2) + \frac{V(t^2)}{z}\right] \tilde{y} \tag{18}$$

$$U(t^2) = X^{-1}(t^2)\ \hat{\mathcal{U}}(t^2)\ X(t^2) = \operatorname{diag}(u_1(t^2), ..., u_k(t^2))$$

$$V(t^2) = X^{-1}(t^2)\ \hat{\mu}\ X(t^2))$$

$$V(t^2)^T + V(t^2) = 0$$

The skew symmetry of V follows from $\eta\hat{\mu} + \hat{\mu}\eta = 0$ and from the choice of the normalization of the eigenvectors, such that $X(t^2)^T\ X(t^2) = \eta^{-1}$.

Let us fix an initial point $t_0 = (0, t_0^2, 0, ..., 0)$. The system (18) has fundamental matrices $y_R(t_0^2, z)$, $y_L(t_0^2, z)$, which are asymptotic to the formal solution

$$\tilde{y}_F(t_0^2, z) = \left[I + \frac{F_1(t_0^2)}{z} + \frac{F_2(t_0^2)}{z} + ...\right]\ e^{z\ U(t_0^2)}$$

in the sectors

$$\mathcal{S}_L(t_0) = \{z \in \mathbf{C} \mid 0 < \arg\left[z \exp\left(\frac{t_0^2}{k}\right)\right] < \pi + \frac{\pi}{k}\}$$

$$\mathcal{S}_R(t_0) = \{z \in \mathbf{C} \mid -\pi < \arg\left[z \exp\left(\frac{t_0^2}{k}\right)\right] < \frac{\pi}{k}\}$$

and

$$\tilde{y}_L(t_0^2, z) = y_R(t_0^2, z)\ S \qquad 0 < \arg\left[z \exp\left(\frac{t_0^2}{k}\right)\right] < \frac{\pi}{k}$$

with respect to the admissible line

$$l_{t_0} := \{z \mid z = \rho \exp\left(i\epsilon - \frac{\Im m\ t_0^2}{k}\right), \quad \rho > 0\}$$

The Stokes' matrix is precisely the matrix S of system (11) with respect to the admissible line l_{t_0}. Also the central connection matrix defined by

$$y_0(t_0^2, z) = y_R(t_0^2, z)\ C \qquad -\pi < \arg\left[ze^{\frac{t_0^2}{k}}\right] < \frac{\pi}{k}$$

is the the same of the system (11).

Definition: C and S, together with $\hat{\mu}$, R, and $e = \frac{\partial}{\partial t^1}$ are the *monodromy data* of the quantum cohomology of $\mathbf{CP}^{k-1}$ in a local chart containing t_0.



Figure 2: Sector where Y_{k2} has the asymptotic behaviour $X\ e^{zU}$ for $z \to \infty$

Recall that we fixed a point $t_0 = (0, t_0^2, ..., 0)$. When we consider a point t away from t_0, the system (17) acquires the general form

$$\partial_z y = \left[\hat{\mathcal{U}}(t) + \frac{\hat{\mu}}{z}\right] y \tag{19}$$

where $\hat{\mathcal{U}}(t^1, .., t^k) = \eta\ \mathcal{U}(t)\ \eta^{-1}$ and $y_\alpha^{(j)}(t^1, .., t^k; z) = \partial_\alpha \tilde{t}^j(t, z)$.

The admissible line l_{t_0} must be considered *fixed once and for all.* Instead, the Stokes' rays change. This is because they are functions of the eigenvalues $u_1(t)$, ..., $u_k(t)$ of the matrix $\hat{\mathcal{U}}(t^1, .., t^k)$. For example, if just t^2 varies, while $t^1 = t^3 = ... = t^k = 0$, the system (17) has Stokes' rays

$$R_{rs}(t^2) = \{z \mid\ = \rho\ \exp\left(i\frac{2\pi}{k} - i\frac{\pi}{k}(r+s) - i\frac{\Im m\ t^2}{k}\right),\quad \rho > 0\}$$

The dependence of the coefficients of the system (19) on t is isomonodromic [9] [10]. Then μ and R are the same for any t. S and C do not change if we move in a sufficently small neighbourhood of t_0. Problems arise when some Stokes' rays cross l_{t_0}. S and C must be modified by an action of the braid group. We will return to this point later.

5 Computation of S

To compute S, we factorize it in "Stokes' factors". Our fundamental matrix Y_L has the required asymptotic form on the sector between R_{k2} $(\arg z = 0)$ and R_{1k} $(\arg z = \pi + \frac{\pi}{k})$. Y_R has the same behaviour between R_{2k} $(\arg z = -\pi)$ and R_{k1} $(\arg z = \frac{\pi}{k})$.

Of course, we can consider fundamental matrices with the same asymptotic behaviour on other sectors of angular width less then $\pi + \frac{\pi}{k}$ and bounded by two Stokes' rays. We introduce the following notation: consider a fundamental matrix of (11) having the required asymptotic behaviour on such a sector. If we go all over the sector clockwise we meet Stokes rays belonging to the sector at each displacement of $\frac{\pi}{k}$. Let R_{ij} be the last ray we meet before reaching the boundary (the boundary is still a Stokes ray *not* belonging to the sector). Then we will call the fundamental matrix Y_{ij}. For example, $Y_L = Y_{k1}$ and $Y_R = Y_{1k}$. See figure 2.

Sometimes, a different labelling is used in the literature. The rays must be enumerated as in figure 3. The numeration refers to the line l: the rays are labelled in counter-clockwise order starting from the first one in Π_R (which will be R_0; then R_0, R_1, ..., R_{k-1} are in Π_R, and R_k, ..., R_{2k-1} are in Π_L). For our particular choiche of l , $R_0 \equiv R_{1k}$ (at $\arg z = -\pi + \frac{\pi}{k}$); $R_1 \equiv R_{1,k-1}$ follows counter-clockwise... Then we proceed untill we reach $R_{k-1} \equiv R_{k2}$ before crossing l, and so on. The fundamental matrices



Figure 3: Sector $\mathcal{S}_k$ and labelling of Stokes' rays

are labelled as we prescribed above, namely Y_j if its sector contains R_j as the last ray met going all over the sector clockwise before the boundary. The sector itself is denoted by $\mathcal{S}_j$. See figure 3.

We define *Stokes' factors* to be the connection matrices K_j such that

$$Y_{j+1}(z) = Y_j(z)K_j$$

on the overlapping region of width π. We warn the reader that also the Stokes' factors will be labelled with both conventions above, according to our convenience (for example $K_0 \equiv K_{1k}$).

As a consequence of the above definitions, we can factorize S as follows

$$Y_L = Y_{k1} = Y_{k2}K_{k2} = Y_{k3}K_{k3}K_{k2} = ...$$

$$... = Y_{1k}K_{1k}K_{1,k-1}K_{k,k-1}K_{k,k-2}...K_{k3}K_{k2} \equiv Y_R S$$

Then

$$S = K_{1k}K_{1,k-1}K_{k,k-1}K_{k,k-2}...K_{k3}K_{k2} \tag{20}$$

We observe that, being the first row of $Y(z)$ equat to $z^{\frac{k-1}{2}}\Phi(z)$, the following holds:

$$\Phi_{j+1}(z) = \Phi_j K_j$$

Remark 3: The Stokes' factors of the system (11) and of the gauge-equivalent system (14) are the same. From the skew-symmetry of V it follows that $K_{ji} = (K_{ij}^{-1})^T$.

Before computing the Stokes factors explicitely, we show that just two of them are enough to compute all the others. Let

$$F(z) := \left(\frac{1}{\sqrt{k}} \frac{1}{z^{\frac{k-1}{2}}} \exp(kz),\ \frac{1}{\sqrt{k}} \frac{e^{\frac{i\pi}{k}}}{z^{\frac{k-1}{2}}} \exp(ke^{\frac{2\pi i}{k}}z),\ ...,\ \frac{1}{\sqrt{k}} \frac{e^{\frac{i\pi}{k}(k-1)}}{z^{\frac{k-1}{2}}} \exp(ke^{\frac{2\pi i}{k}(k-1)}z) \right)$$

$$= (F^{(1)}(z),\ F^{(2)}(z),\ ...,\ F^{(k)}(z))$$

be the row vector whose entries are the first terms of the asymptotic expansions of an actual solution $\Phi(z)$ of the generalized hypergeometric equation. By a straightforward computation we see that

$$F(ze^{\frac{2\pi i}{k}}) = F(z)T_F\ , \qquad T_F = \begin{pmatrix} 0 & & & & ... & 1 \\ -1 & 0 & & & & \\ & -1 & 0 & & & \\ & & -1 & & & \\ & & & \ddots & \ddots & \vdots \\ & & & & -1 & 0 \end{pmatrix}$$

We use now the convention of enumeration of Stokes' rays R_0, R_1, ... starting from l (see above). Let $\Phi_m(z)$ be an actual solution of the hypergeometric equation having asymptotic behaviour $F(z)$ on $\mathcal{S}_m$.

$$\Phi_m(z) \sim F(z) \quad z \to \infty \quad z \in \mathcal{S}_m$$

Then

$$\Phi_{m+2}(ze^{\frac{2\pi i}{k}}) \sim F(ze^{\frac{2\pi i}{k}}) = F(z) T_F \ , \qquad z \in \mathcal{S}_m$$

Namely,

$$\Phi_{m+2}(ze^{\frac{2\pi i}{k}}) T_F^{-1} \sim F(z) \ , \qquad z \in \mathcal{S}_m$$

then, by unicity of actual solutions having asymptotic behaviour $F(z)$ in a sector wider then π, we have

$$\Phi_{m+2}(ze^{\frac{2\pi i}{k}}) = \Phi_m(z) T_F \ , \qquad z \in \mathcal{S}_m$$

Lemma 6: *For any* $p \in \mathbf{Z}$

$$K_{m+2p} = T_F^{-p} K_m T_F^p$$

Proof: For $z \in \mathcal{S}_m \cap \mathcal{S}_{m+1}$

$$\Phi_{m+1}(z) = \Phi_m(z) \ K_m = \Phi_{m+2}(ze^{\frac{2\pi i}{k}}) \ T_F^{-1} K_m =$$

$$= \Phi_{m+3}(ze^{\frac{2\pi i}{k}}) \ K_{m+2}^{-1} T_F^{-1} K_m = \Phi_{m+1}(z) \ \ T_F \ K_{m+2}^{-1} \ T_F^{-1} \ K_m$$

Then $K_{m+2} = \ T_F^{-1} \ K_m \ T_F$. By induction we prove the lemma.

□

¿From the lemma it follows that just two Stokes' factors are enought to compute all the others. We are ready to give a concise formula for S:

Theorem 1: *Let l be an admissible line (i.e. not containing Stokes' rays), and let us enumerate the rays in counter-clockwise order starting from the first one in Π_R (which will be R_0, then R_0, R_1, ..., R_{k-1} are in Π_R, and R_k, ..., R_{2k-1} are in Π_L). Then the Stokes' matrix for (11), (14), (13) and $k > 3$ is*

$$S = \begin{cases} \left(K_0 \ K_1 \ T_F^{-1}\right)^{\frac{k}{2}} \ T_F^{\frac{k}{2}} \ \equiv \ T_F^{\frac{k}{2}} \ \left(T_F^{-1} \ K_{k-2} \ K_{k-1}\right)^{\frac{k}{2}}, & k \ \text{even} \\ \\ \left(K_0 \ K_1 \ T_F^{-1}\right)^{\frac{k-1}{2}} \ K_0 \ T_F^{\frac{k-1}{2}} \ \equiv \ T_F^{\frac{k-1}{2}} \ K_{k-1} \left(T_F^{-1} \ K_{k-2} \ K_{k-1}\right)^{\frac{k-1}{2}}, & k \ \text{odd} \end{cases} \tag{21}$$

Proof: $S = K_0 \ K_1 \ K_2 \ ... \ K_{k-1}$, $K_{2p} = T_F^{-p} \ K_0 \ T_F^p$ and $K_{2p+1} = T_F^{-p} \ K_1 \ T_F^p$. Then

$$S = K_0 K_1 \ (T_F^{-1} K_0 T_F) \ (T_F^{-1} K_1 T_F) \ K_4 K_5 \ ... \ K_{k-1}$$

$$= (K_0 K_1 T_F^{-1}) \ K_0 K_1 \ T_F \ K_4 K_5 \ ... \ K_{k-1}$$

$$= (K_0 K_1 T_F^{-1}) \ K_0 K_1 \ T_F \ (T_F^{-2} K_0 T_F^2) \ (T_F^{-2} K_1 T_F^2) \ K_6 \ ... \ K_{k-1}$$

$$= (K_0 K_1 T_F^{-1})(K_0 K_1 T_F^{-1}) \ K_0 K_1 T_F^2 \ K_6 \ ... \ K_{k-1}$$

Now observe that $K_{k-1} = T_F^{-(\frac{k}{2}-1)} \ K_1 \ T_F^{\frac{k}{2}-1}$ for k even, while $K_{k-1} = T_F^{-(\frac{k-1}{2})} \ K_0 \ T_F^{\frac{k-1}{2}}$ for k odd. Then, for k even

$$S = (K_0 K_1 T_F^{-1})^{\frac{k}{2}-2} \ K_0 K_1 T_F^{\frac{k}{2}-2} \ K_{k-2} K_{k-1}$$

$$= (K_0 K_1 T_F^{-1})^{\frac{k}{2}-2} \ K_0 K_1 T_F^{\frac{k}{2}-2} \ T_F^{-(\frac{k}{2}-1)} K_0 T_F^{\frac{k}{2}-1} \ T_F^{-(\frac{k}{2}-1)} K_1 T_F^{\frac{k}{2}-1} = \left(K_0 \ K_1 \ T_F^{-1}\right)^{\frac{k}{2}} \ T_F^{\frac{k}{2}}$$

For k odd:

$$S = (K_0 K_1 T_F^{-1})^{\frac{k-3}{2}} \ K_0 K_1 T_F^{\frac{k-3}{2}} \ K_{k-1}$$

$$= (K_0 K_1 T_F^{-1})^{\frac{k-3}{2}} \ K_0 K_1 T_F^{\frac{k-3}{2}} \ T_F^{-(\frac{k-1}{2})} K_0 T_F^{\frac{k-1}{2}} = \left(K_0 \ K_1 \ T_F^{-1}\right)^{\frac{k-1}{2}} \ K_0 \ T_F^{\frac{k-1}{2}}$$

If instead we write the Stokes' factors in term of K_{k-2} and K_{k-1} we obtain the other two formulas in the same way.

$\square$

Remark 4: For our particular choiche of l, $K_0 \equiv K_{1k}$, $K_1 \equiv K_{1,k-1}$, $K_{k-2} \equiv K_{k3}$ and $K_{k-1} \equiv K_{k2}$. For $k = 3$

$$S = T_F K_{32}\ (T_F^{-1} K_{12} K_{32}) = K_{13} K_{12} K_{32}$$

It is now worth deriving some properties of the monodromy of $Y(z)$ (for (11)) and $\Phi(z)$ (for (13)), which will be usefull later. Consider $\Phi_m(z)$ with asymptotic behaviour $F(z)$ on $\mathcal{S}_m$. Then

$$\Phi_m(z) = \Phi_{m-2}(z)\ K_{m-2}\ K_{m-1} \equiv \Phi_m(ze^{\frac{2\pi i}{k}})\ T_F^{-1}\ K_{m-2}\ K_{m-1}$$

On the other hand

$$\Phi_m(z) = \Phi_{m+2}(z)\ K_{m+1}^{-1}\ K_m^{-1} \equiv \Phi_m(ze^{-\frac{2\pi i}{k}})\ T_F\ K_{m+1}^{-1}\ K_m^{-1}$$

This proves the following

Lemma 7: *The basic solution $\Phi_m(z)$ of the generalized hypergeometric equation (13) with asymptotic behaviour $F(z)$ on $\mathcal{S}_m$, satisfies the identity*

$$\Phi_m(ze^{\frac{2\pi i}{k}}) = \Phi_m(z)\ T_m$$

where

$$T_m := \ K_{m-1}^{-1}\ K_{m-2}^{-1}\ T_F\ \ =\ T_F\ K_{m+1}^{-1}\ K_m^{-1}$$

Corollary 1: *The monodromy (at $z = 0$) of $\Phi_m(z)$ is*

$$\Phi_m(ze^{2\pi i}) = \Phi_m(z)\ (T_m)^k$$

Now, for our particular choiche of the line l and for $m = k$, $\Phi_m(z) = \Phi_L(z)$. For the solution $Y_L(z)$ of (11), the relations $Y_R(z) = Y_L(z)\ S^{-1}$ $(0 < \arg z < \frac{\pi}{k})$, $Y_L(z) = Y_R(ze^{-2\pi i})\ S^T$ $(\pi < \arg z < \pi + \frac{\pi}{k})$ immediately imply

$$Y_L(ze^{2\pi i}) = Y_L(z)\ S^{-1}\ S^T$$

Recall that the (n, j)-th entry of $Y(z)$ is $Y_{n,j}(z) \equiv Y_n^{(j)}(z) = \frac{1}{k^{n-1}}\ z^{\frac{k-1}{2}-n+1}\ (z\partial_z)^{(n-1)}\ \varphi^{(j)}(z)$, and observe that $(ze^{2\pi i})^{\frac{k-1}{2}} = (-1)^{k-1}\ z^{\frac{k-1}{2}}$ Then, from Corollary 1 we get the following:

Corollary 2: *Let T be the k-monodromy matrix of Φ_L (namely, $T \equiv T_k$ for our choiche of l). Then*

$$T^k = (-1)^{k-1}\ S^{-1}\ S^T$$

Our formula (21) allows us to easily compute S. The recipe is simply to take K_{k3}, K_{k2} (which we are going to compute explicitely) and substitute them into

$$S = T_F^{\frac{k}{2}}\ (T_F^{-1}\ K_{k3}\ K_{k2})^{\frac{k}{2}} = T_F^{\frac{k}{2}}\ T^{-\frac{k}{2}} \tag{22}$$

for k even, or into

$$S = T_F^{\frac{k-1}{2}}\ K_{k2}\ (T_F^{-1}\ K_{k3}\ K_{k2})^{\frac{k-1}{2}} = T_F^{\frac{k-1}{2}}\ K_{k2}\ T^{-\frac{k-1}{2}} \tag{23}$$

for k odd.

Computation of Stokes' factors: We need to distinguish between k odd and even. In the following $g(z)$ will mean $g^{(n(k))}(z)$ $(n(k) = \frac{k}{2} + 1$ or $\frac{k+1}{2}$ for k even or odd respectively).

k odd:

$$g(z) = \varphi_L^{(\frac{k+1}{2})}(z) \equiv \varphi_R^{(\frac{k+1}{2})}(z)$$

with asymptotic behaviour on

$$-\pi < \arg z < \pi + \frac{2\pi}{k}$$

If we iterate the map $z \mapsto ze^{\frac{2\pi i}{k}}$ for $m = 1, 2, ..., \frac{k+1}{2}$ times, the domain of $g(ze^{\frac{2\pi i}{k}m})$ for each m covers $\mathcal{S}_R$. When we reach $m = \frac{k+1}{2}$ a new iteration (i.e. a new rotation of the domain of $-\frac{2\pi}{k}$) will live the sector $-\frac{\pi}{k} < \arg z < \frac{\pi}{k}$ of $\mathcal{S}_R$ uncovered. The same, if we do $z \mapsto ze^{-\frac{2\pi i}{k}}$ the sector $-\pi < \arg z < -\pi + \frac{2\pi}{k}$ of $\mathcal{S}_R$ remains uncovered. Then, by remark 2:

$$\Phi_{1k}(z) \equiv \Phi_R(z) = \Big((-1)^{\frac{k-1}{2}} g(ze^{i\frac{2\pi}{k}(\frac{k+1}{2})}),\ ...\ \tfrac{k-3}{2} \text{ unknown terms } ...\ ,$$

$$g(z),\ -g(ze^{\frac{2\pi i}{k}}),\ g(ze^{\frac{4\pi i}{k}}),\ ...\ ,\ (-1)^{\frac{k-1}{2}} g(ze^{i\frac{2\pi}{k}(\frac{k-1}{2})})\Big)$$

In the same way we see that

$$\Phi_{k1}(z) \equiv \Phi_L(z) = \Big((-1)^{\frac{k-1}{2}} g(ze^{-i\frac{2\pi}{k}(\frac{k-1}{2})}),\ -(-1)^{\frac{k-1}{2}} g(ze^{-i\frac{2\pi}{k}(\frac{k-3}{2})}),\ ...\ , -g(ze^{-\frac{2\pi i}{k}}),$$

$$g(z),\ ...\ \tfrac{k-1}{2} \text{ unknown terms } ...\Big)$$

and similar expressions for $\Phi_{1,k-1},\ \Phi_{k,k-1},\ \Phi_{k,k-2},\ ...,\ \Phi_{k3}, \Phi_{k2}$. The unknown terms are computed using the identity (16) and simple considerations on the dominant exponentials $|e^{zu_i}|$ on the sectors which remain uncovered in the iterations of $z \mapsto ze^{\pm i\frac{2\pi}{k}}$.

Example: A simple example will clarify this procedure. Let $k = 7$; then $g = \varphi^{(4)}$,

$$\Phi_{17} = \Phi_R = (-g(ze^{\frac{8\pi i}{7}}),\ ?,\ ?,\ g(z),\ -g(ze^{\frac{2\pi i}{7}}),\ g(ze^{\frac{4\pi i}{7}}),\ -g(ze^{\frac{6\pi i}{7}}))$$

We look for $\varphi_R^{(3)}$. If we take $(-1)g(ze^{\frac{-2\pi i}{k}})$ we miss to cover $-\pi < \arg z < -\pi + \frac{2\pi}{7}$ in $\mathcal{S}_R$. On $-\pi < \arg z < -\pi + \frac{\pi}{7}$ (between two nearby Stokes' rays) we have the relations $|e^{zu_4}| > |e^{zu_5}| > |e^{zu_3}|$. On $-\pi + \frac{\pi}{7} < \arg z < -\pi + \frac{2\pi}{7}$, $|e^{zu_4}| > |e^{zu_3}|$ (later on we will simply write $4 > 3$). Then

$$\varphi_R^{(3)}(z) = (-1)g(ze^{\frac{-2\pi i}{7}}) + c_4\varphi_R^{(4)}(z) + c_5\varphi_R^{(5)}(z)$$

To find c_4, c_5 we need another representation for $\varphi_R^{(3)}$. We consider $(-1)g(ze^{\frac{12\pi i}{7}})$, which has the correct asymptotic behaviour, but on a domain which leaves uncovered $-\frac{3\pi}{7} < \arg z < \frac{\pi}{7}$. The relations are: on $0 < \arg z < \frac{\pi}{7}$, $1 > 7 > 2 > 6 > 3$; on $-\frac{\pi}{7} < \arg z < 0$, $1 > 2 > 7 > 3$; on $-\frac{2\pi}{7} < \arg z < -\frac{\pi}{7}$, $2 > 1 > 3$; on $-\frac{3\pi}{7} < \arg z < -\frac{2\pi}{7}$, $2 > 3$. Then

$$\varphi_R^{(3)}(z) = (-1)g(ze^{\frac{12\pi i}{7}}) + d_1\varphi_R^{(1)}(z) + d_2\varphi_R^{(2)}(z) + d_6\varphi_R^{(6)}(z) + d_7\varphi_R^{(7)}(z)$$

In the same way one finds

$$\varphi_R^{(2)}(z) = \begin{cases} g(ze^{\frac{-4\pi i}{7}}) + a_3\varphi_R^{(3)}(z) + a_4\varphi_R^{(4)}(z) + a_5\varphi_R^{(5)}(z) + a_6\varphi_R^{(6)}(z) \\ g(ze^{\frac{10\pi i}{7}}) + b_1\varphi_R^{(1)}(z) + b_2\varphi_R^{(4)}(z) \end{cases}$$

$\varphi^{(i)}$'s are known for $i = 1, 4, 5, 6, 7$. Using the identity (16) we compute a, b, c, d. We get

$$\varphi_R^{(2)}(z) = g(ze^{\frac{-4\pi i}{7}}) - \binom{7}{1} g(ze^{\frac{-2\pi i}{7}}) + \binom{7}{2} g(z) - \binom{7}{3} g(ze^{\frac{2\pi i}{7}}) + \binom{7}{4} g(ze^{\frac{4\pi i}{7}})$$

$$\varphi_R^{(3)}(z) = -g(ze^{\frac{-2\pi i}{7}}) + \binom{7}{1} g(z) - \binom{7}{2} g(ze^{\frac{2\pi i}{7}})$$

A similar computation gives $\Phi_{71} = \Phi_L$.

$$\Phi_{71}^T = \begin{bmatrix} -g(ze^{-\frac{6\pi i}{7}}) \\ g(ze^{-\frac{4\pi i}{7}}) \\ -g(ze^{\frac{-2\pi i}{7}}) \\ g(z) \\ -g(ze^{\frac{2\pi i}{7}}) + \binom{7}{6} g(z) \\ g(ze^{\frac{4\pi i}{7}}) - \binom{7}{6} g(ze^{\frac{2\pi i}{7}}) + \binom{7}{5} g(z) - \binom{7}{4} g(ze^{-\frac{2\pi i}{7}}) \\ -g(ze^{\frac{6\pi i}{7}}) + \binom{7}{6} g(ze^{\frac{4\pi i}{7}}) - \binom{7}{5} g(ze^{\frac{2\pi i}{7}}) + \binom{7}{4} g(z) - \binom{7}{3} g(ze^{-\frac{2\pi i}{7}}) + \binom{7}{2} g(ze^{-\frac{4\pi i}{7}}) \end{bmatrix}$$

and

$$\Phi_{72}^T = \begin{bmatrix} -g(ze^{-\frac{6\pi i}{7}}) \\ g(ze^{-\frac{4\pi i}{7}}) \\ -g(ze^{\frac{-2\pi i}{7}}) \\ g(z) \\ -g(ze^{\frac{2\pi i}{7}}) \\ g(ze^{\frac{4\pi i}{7}}) - \begin{pmatrix} 7 \\ 6 \end{pmatrix} g(ze^{\frac{2\pi i}{7}}) + \begin{pmatrix} 7 \\ 5 \end{pmatrix} g(z) \\ -g(ze^{\frac{6\pi i}{7}}) + \begin{pmatrix} 7 \\ 6 \end{pmatrix} g(ze^{\frac{4\pi i}{7}}) - \begin{pmatrix} 7 \\ 5 \end{pmatrix} g(ze^{\frac{2\pi i}{7}}) + \begin{pmatrix} 7 \\ 4 \end{pmatrix} g(z) - \begin{pmatrix} 7 \\ 3 \end{pmatrix} g(ze^{-\frac{2\pi i}{7}}) \end{bmatrix}$$

Notice that in each of the last three entries of Φ_{72} there is a term missing w.r.t the corresponding entries of Φ_{71}. This immediately implies

$$K_{72} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & \binom{7}{2} \\ 0 & 0 & 1 & 0 & 0 & \binom{7}{4} & 0 \\ 0 & 0 & 0 & 1 & \binom{7}{6} & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The next step is the computation of Φ_{73} and K_{73}, through $\Phi_{72} = \Phi_{73} K_{73}$. It is done in the same way...

The above procedure is extended to the general case. In Appendix 2 we give, for example, the general expressions of Φ_R and Φ_L. The factors of interest are:

k **odd:**

$(K_{k2})_{j,\ k-j+2} = \begin{pmatrix} k \\ 2(j-1) \end{pmatrix}$ for $j = 2, ..., \frac{k+1}{2}$. $(K_{k2})_{j,j} = 1$ for $j = 1, ..., k$. All the other entries are zero.

$(K_{k3})_{2,1} = -\begin{pmatrix} k \\ 1 \end{pmatrix}$; $(K_{k3})_{j,\ k-j+3} = \begin{pmatrix} k \\ 2j-3 \end{pmatrix}$ for $j = 3, ..., \frac{k+1}{2}$. $(K_{k3})_{j,j} = 1$ for $j = 1, ..., k$. All the other entries are zero. Namely:

$$K_{k2} = \begin{pmatrix} 1 & & & & & & & & & & 0 \\ & 1 & & & & & & & & & \binom{k}{2} \\ & & 1 & & & & & & & \binom{k}{4} & \\ & & & 1 & & & & & \binom{k}{6} & & \\ & & & & \ddots & & & \cdot^{\cdot^{\cdot}} & & & \\ & & & & & 1 & \binom{k}{k-1} & & & & \\ & & & & & & 1 & & & & \\ & & & & & & & \ddots & & & \\ & & & & & & & & 1 & & \\ & & & & & & & & & 1 & \\ & & & & & & & & & & 1 \end{pmatrix}$$

$$K_{k3} = \begin{pmatrix} 1 & & & & & & & & & & & & 0 \\ -\binom{k}{1} & 1 & & & & & & & & & & & 0 \\ & & 1 & & & & & & & & & & \binom{k}{3} \\ & & & 1 & & & & & & & & \binom{k}{5} & \\ & & & & 1 & & & & & & \binom{k}{7} & & \\ & & & & & \ddots & & & & \cdot^{\cdot^{\cdot}} & & & \\ & & & & & & 1 & 0 & \binom{k}{k-2} & & & & \\ & & & & & & & 1 & 0 & & & & \\ & & & & & & & & 1 & & & & \\ & & & & & & & & & \ddots & & & \\ & & & & & & & & & & 1 & & \\ & & & & & & & & & & & 1 & \\ & & & & & & & & & & & & 1 \end{pmatrix}$$

k **even:**

$(K_{k2})_{j,\ k-j+2}=\begin{pmatrix}k\\ 2(j-1)\end{pmatrix}$ for $j=2,...,\frac{k}{2}$. $(K_{k2})_{j,j}=1$ for $j=1,...,k$. All the other entries are zero.

$(K_{k3})_{2,1}=-\begin{pmatrix}k\\ 1\end{pmatrix}$; $(K_{k3})_{j,\ k-j+3}=\begin{pmatrix}k\\ 2j-3\end{pmatrix}$, for $j=3,...,\frac{k}{2}+1$. $(K_{k3})_{j,j}=1$ for $j=1,...,k$. All the other entries are zero. Namely

$$K_{k2}=\begin{pmatrix}
1 & & & & & & & & & & & 0\\
 & 1 & & & & & & & & & & \binom{k}{2}\\
 & & 1 & & & & & & & & \binom{k}{4} & \\
 & & & 1 & & & & & & \binom{k}{6} & & \\
 & & & & \ddots & & & & \cdot^{\cdot^{\cdot}} & & & \\
 & & & & & 1 & 0 & \binom{k}{k-2} & & & & \\
 & & & & & & 1 & 0 & & & & \\
 & & & & & & & 1 & & & & \\
 & & & & & & & & \ddots & & & \\
 & & & & & & & & & 1 & & \\
 & & & & & & & & & & 1 & \\
 & & & & & & & & & & & 1
\end{pmatrix}$$

$$K_{k3}=\begin{pmatrix}
1 & & & & & & & & & & & & 0\\
-\binom{k}{1} & 1 & & & & & & & & & & & 0\\
 & & 1 & & & & & & & & & & \binom{k}{3}\\
 & & & 1 & & & & & & & & \binom{k}{5} & \\
 & & & & 1 & & & & & & \binom{k}{7} & & \\
 & & & & & \ddots & & & & \cdot^{\cdot^{\cdot}} & & & \\
 & & & & & & 1 & \binom{k}{k-1} & & & & & \\
 & & & & & & & 1 & & & & & \\
 & & & & & & & & & \ddots & & & \\
 & & & & & & & & & & 1 & & \\
 & & & & & & & & & & & 1 & \\
 & & & & & & & & & & & & 1
\end{pmatrix}$$

6 Reduction of S to "Canonical" Form

Some examples of computations of S are in Appendix 2. The reader may observe that S is not in a nice upper triangular form (see also Lemma 4) and quite strange numbers (complicated combinations of sum and products of binomial coefficients) appear.

Some natural operations are allowed on the Stokes matrices of a Frobenius manifold:

a) Permutations. Let us consider the system

$$\frac{d\tilde{Y}}{dz}=\left[U+\frac{1}{z}V\right]\ \tilde{Y}$$

where $U=\mathrm{diag}(u_1,\ u_2,\ ...,\ u_k)$. Let $\sigma:\ (1,2,..,k)\mapsto(\sigma(1),\sigma(2),...,\sigma(k))$ be a permutation. It is represented by an invertible matrix P such that

$$P\ U\ P^{-1}\ =\ \mathrm{diag}(u_{\sigma(1)},u_{\sigma(2)},...,u_{\sigma(k)})$$

The system

$$\frac{d\mathcal{Y}}{dz}=\left[PUP^{-1}+\frac{1}{z}PVP^{-1}\right]\ \mathcal{Y}$$

has solutions

$$\mathcal{Y}_{L/R}(z):=P\ \tilde{Y}_{L/R}(z)\ P^{-1}\sim(I+O(\frac{1}{z}))\ e^{zPUP^{-1}}\qquad \mathcal{Y}_L(z)=\mathcal{Y}_R(z)\ PSP^{-1}$$

$$\mathcal{Y}_0(z):=P\tilde{Y}_0(z)=(PX^{-1}+O(z))z^{\mu}z^{R},\quad \mathcal{Y}_0(z)=\mathcal{Y}_R(z)\ PC$$

S and C are then transformed in PSP^{-1} and PC. For a suitable P, PSP^{-1} is upper triangular. As a general result [1], the good permutation is the one which put $u_{\sigma(1)}$, ..., $u_{\sigma(k)}$ in lexicographical ordering w.r.t. the oriented line l. The effect of the permutation P corresponds to a change of coordinates in the given chart, consisting in the permutation σ of the coordinates.

b) Sign changes of the entries. The construction at point a) is repeated, but now a diagonal matrix $\mathcal{I}$ with 1's or -1's on the diagonal takes the place of P. In $\mathcal{I}S\mathcal{I}^{-1}$ some entries change sign. Note that $\mathcal{I}U\mathcal{I}^{-1} \equiv U$. Moreover, the formulae [9] [10] which define a local chart of the manifold in terms of monodromy data are not affected by $S \mapsto \mathcal{I}S\mathcal{I}$, $C \mapsto \mathcal{I}C$.

c) Action of the braid group.
We first recall that the braid group is generated by $k-1$ elementary braids β_{12}, β_{23}, ..., $\beta_{k-1,k}$, with relations:

$$\beta_{i,i+1}\beta_{j,j+1} = \beta_{j,j+1}\beta_{i,i+1} \quad \text{for } i+1 \neq j, \; j+1 \neq i$$

$$\beta_{i,i+1}\beta_{i+1,i+2}\beta_{i,i+1} = \beta_{i+1,i+2}\beta_{i,i+1}\beta_{i+1,i+2}$$

This abstract group is realized as the fundamental group of $(\mathbf{C}^k \backslash \text{diagonals})/\mathcal{S}_k := \{(u_1, ..., u_k) \mid u_i \neq u_j \text{ for } i \neq j\}/\mathcal{S}_k$, where $\mathcal{S}_k$ is the symmetric group of order k.

In section 4 we proved that the Stokes' rays of the systems (17) (18), and more generally the rays of the system (19), depend on the point t of the manifold. This is equivalent to the fact that the eigenvalues $u_1(t)$, ..., $u_k(t)$ of $\hat{\mathcal{U}}(t)$ depend on t. When t changes, $u_1(t)$, ..., $u_k(t)$ change their position in the complex plane. Consequently, Stokes' rays move. Let us start from the point $t_0 = (0, t_0^2, ..., 0)$. If we move in a sufficently small neighbourhood of t_0, the rays slightly change their positions. But if we move sufficently far away from t_0, some Stokes' rays cross the *fixed* admissible line l_{t_0}. Then, we must change "Left" and "Right" solutions of (19). Then also S and C change. This is the reason why the monodromy data S and C change when we go from one local chart to another.

The motions of the points $u_1(t)$, ..., $u_k(t)$ as t changes represent transformations of the braid group. Actually, a braid $\beta_{i,i+1}$ can be represented as an "elementary" deformation consisting of a permutation of u_i, u_{i+1} moving counter-clockwise (clockwise or counter-clockwise is a matter of convention).

Suppose u_1, ..., u_k are already in lexicographical order w.r.t. l, so that S is upper triangular (recall that this configuration can be reached by a suitable permutation P). The effect on S of the deformation of u_i, u_{i+1} representing $\beta_{i,i+1}$ is the following :

$$S \mapsto \; S^{\beta_{i,i+1}} := A^{\beta_{i,i+1}}(S) \; S \; A^{\beta_{i,i+1}}(S)$$

where

$$\left(A^{\beta_{i,i+1}}(S)\right)_{nn} = 1 \qquad n = 1, ..., k \quad n \neq \; i, \; i+1$$

$$\left(A^{\beta_{i,i+1}}(S)\right)_{i+1,i+1} = -s_{i,i+1}$$

$$\left(A^{\beta_{i,i+1}}(S)\right)_{i,i+1} = \left(A^{\beta_{i,i+1}}(S)\right)_{i+1,i} = 1$$

and all the other entries are zero.

For the inverse braid $\beta_{i,i+1}^{-1}$ (u_i and u_{i+1} move clockwise) the representation is

$$\left(A^{\beta_{i,i+1}^{-1}}(S)\right)_{nn} = 1 \qquad n = 1, ..., k \quad n \neq \; i, \; i+1$$

$$\left(A^{\beta_{i,i+1}^{-1}}(S)\right)_{i,i} = -s_{i,i+1}$$

$$\left(A^{\beta_{i,i+1}^{-1}}(S)\right)_{i,i+1} = \left(A^{\beta_{i,i+1}^{-1}}(S)\right)_{i+1,i} = 1$$

and all the other entries are zero.

For a generic braid β, which is a product of N elementary braids (for some N) $\beta = \beta_{j_1,j_1+1} ... \beta_{j_N,j_N+1}$, the action is

$$S \mapsto S^{\beta} = A^{\beta}(S) \; S \; \left[A^{\beta}(S)\right]^T$$

where

$$A^{\beta}(S) = A^{\beta_{i_N,i_N+1}}(S^{\beta_{j_1,j_1+1} ... \beta_{j_{N-1},j_{N-1}+1}}) \; ... \; A^{\beta_{i_2,i_2+1}}(S^{\beta_{i_1,i_1+1}}) A^{\beta_{i_1,i_1+1}}(S)$$

We remark that S^{β} is still upper triangular.



Figure 4: Effect of the braid which brings S to the canonical form on the lines L_j

In figure 4 we have drawn some lines $L_j = \{\lambda = u_j + \rho e^{i(\frac{\pi}{2}-\epsilon)}, \quad \rho > 0\}$ ($0 < \epsilon < \frac{\pi}{k}$ is the angle of l), which help us to visualize the topological effect of the braids action (they are the branch cuts for the fuchsian system which will be introduced in section 8). We are going to prove that the braid whose effect is to set the deformed points in cyclic order and the cuts in the configuration of figure 4 (namely, the last two cuts remain unchanged, the others are alternatively "inverted"), brings S in a *canonical form*: $s_{i,i+1} = \begin{pmatrix} k \\ 1 \end{pmatrix}$ $\forall i = 1, ..., k-2$; $s_{i,i+2} = \begin{pmatrix} k \\ 2 \end{pmatrix}$ $\forall i = 1, ..., k-3$; ...; $s_{i,i+n} = \begin{pmatrix} k \\ n \end{pmatrix}$ $\forall i = 1, ..., k-n+1$; $s_{i,k} = -\begin{pmatrix} k \\ k-i \end{pmatrix}$. Namely

$$S^\beta = \begin{pmatrix} 1 & \begin{pmatrix} k \\ 1 \end{pmatrix} & \begin{pmatrix} k \\ 2 \end{pmatrix} & \begin{pmatrix} k \\ 3 \end{pmatrix} & \begin{pmatrix} k \\ 4 \end{pmatrix} & ... & -\begin{pmatrix} k \\ k-1 \end{pmatrix} \\ & 1 & \begin{pmatrix} k \\ 1 \end{pmatrix} & \begin{pmatrix} k \\ 2 \end{pmatrix} & \begin{pmatrix} k \\ 3 \end{pmatrix} & ... & -\begin{pmatrix} k \\ k-2 \end{pmatrix} \\ & & 1 & \begin{pmatrix} k \\ 1 \end{pmatrix} & \begin{pmatrix} k \\ 2 \end{pmatrix} & ... & -* \\ & & & 1 & \begin{pmatrix} k \\ 1 \end{pmatrix} & ... & -* \\ & & & & 1 & & -* \\ & & & & & \ddots & \\ & & & & & & 1 \end{pmatrix} \tag{24}$$

Note that the last column is negative. Its sign is inverted by $S \mapsto \mathcal{I}S\mathcal{I}$, where $\mathcal{I} := \text{diag}(1, 1, ..., -1)$.

Lemma 8: *Let the points u_j ($j = 1, .., k$) be in lexicographical order w.r.t the admissible line l. Then*



Figure 5: Effect of the sequences of braids which bring S to the canonical form (the figure refers to k even)

the braid

$$\beta := (\beta_{k-5,k-4}\ \beta_{k-6,k-5}\ \ldots\ \beta_{12})\ (\beta_{k-6,k-5}\ \beta_{k-7,k-6}\ \ldots\ \beta_{23})\ (\beta_{k-7,k-6}\ \ldots\ \beta_{34})\ \ldots$$

$$\ldots\ \beta_{\frac{k}{2}-2,\frac{k}{2}-1}\ (\beta_{k-3,k-2}\ \beta_{k-4,k-3}\ \ldots\ \beta_{12})$$

for k even, or

$$\beta := (\beta_{k-5,k-4}\ \beta_{k-6,k-5}\ \ldots\ \beta_{12})\ (\beta_{k-6,k-5}\ \beta_{k-7,k-6}\ \ldots\ \beta_{23})\ (\beta_{k-7,k-6}\ \ldots\ \beta_{34})\ \ldots$$

$$\ldots\ (\beta_{\frac{k-3}{2},\frac{k-1}{2}}\ \beta_{\frac{k-5}{2},\frac{k-3}{2}})\ (\beta_{k-3,k-2}\ \beta_{k-4,k-3}\ \ldots\ \beta_{12})$$

for k odd, brings the points in cyclic counter-clockwise order, u_1 being the first point in Π_L (figure 4, right side, or figure 5).

Note that we have collected the braids in $\frac{k}{2}-1$ (k even), or $\frac{k-3}{2}$ (k odd) sequences (...).

Proof: Let k be even. The first braid $\beta_{k-5,k-4}$ iterchanges u_{k-4} and u_{k-5}. The second braid interchanges u_{k-5} and u_{k-6}. One easily sees that the effect of the first sequence of braids $(\beta_{k-5,k-4}\ \beta_{k-6,k-5}\ \ldots\ \beta_{12})$ is to bring u_1 in the (old) position of u_{k-4}, u_{k-4} in the position of u_{k-5}, u_{k-5} in the position of u_{k-6}, ..., u_4 in the position of u_3 and u_2 in the position of u_1 (figure 5). u_k, u_{k-1}, u_{k-2}, u_{k-3} are not moved.

The second sequence of braids $(\beta_{k-6,k-5}\ \beta_{k-7,k-6}\ \ldots\ \beta_{23})$ acts in a similar way, bringing u_2 in u_{k-5}, u_{k-5} in u_{k-6}, ..., u_3 in u_2. u_k, u_{k-1}, u_{k-2}, u_{k-3}, u_{k-4} are not moved.

We go on in this way. After the action of

$$(\beta_{k-5,k-4}\ \beta_{k-6,k-5}\ \ldots\ \beta_{12})\ (\beta_{k-6,k-5}\ \beta_{k-7,k-6}\ \ldots\ \beta_{23})\ (\beta_{k-7,k-6}\ \ldots\ \beta_{34})\ \ldots\ \beta_{\frac{k}{2}-2,\frac{k}{2}-1}$$

the points are as in figure 5: u_k is on the positive real axis, u_{k-2} is the first point met in counter-clockwise order, u_1 is the second, u_2 is the third; the points are in cyclic order up to u_{k-3}; finally, u_{k-1} is the last point before reaching again the positive real axis from below.

Then, $(\beta_{k-3,k-2}\ \beta_{k-4,k-3}\ \ldots\ \beta_{12})$ brings u_1 in u_{k-2}, u_{k-2} in u_{k-3}, u_{k-3} in u_{k-4}, and so on. The cyclic order is reached.

For k odd the proof is similar.

□

A careful consideration of the effect of the braid β on the lines L_j (which we leave as an exercise for the reader) shows that they are alternatively inverted as in figure 4. To reconstruct uniquely this configuration we just need to know the oriented line l, namely, its angle ϵ w.r.t the positive real axis. The points u_{k-1}, u_k and the lines L_{k-1} and L_k are unchanged (angle $\frac{\pi}{2}-\epsilon$). The line at u_1 starts in the opposite direction, it goes around $u_2, \ldots, u_{k-2}$ without intesecting other cuts, and then goes to ∞ with the original asymptotic direction $\frac{\pi}{2}-\epsilon$. Moving in the direction opposite to that of l we meet u_{k-2}. Its line has the original direction $\frac{\pi}{2}-\epsilon$. Then we meet u_2, and the corresponding line starts with opposite direction, goes arouns $u_3, \ldots, u_{k-3}$ and then goes to ∞ with asymptotic direction $\frac{\pi}{2}-\epsilon$. And so on.

Now we find the matrix representation for the braid β.

Proposition 1: *The braid β of Lemma 8 has the following matrix representation:*

$$A^{\beta}(S)=\begin{pmatrix}
0 & 0 & 0 & 0 & 0 & & 0 & 0 & 0 & 0 & 1 & 0 & 0\\
 & & & & & & 0 & 0 & 1 & 0 & \binom{k}{1} & 0 & 0\\
 & & & & & & 1 & 0 & \binom{k}{1} & 0 & \binom{k}{2} & 0 & 0\\
 & & & & & & \binom{k}{1} & 0 & \binom{k}{2} & 0 & \binom{k}{3} & 0 & 0\\
 & & & & & & * & . & * & . & * & . & .\\
 & & & & & \cdots & * & . & * & . & * & . & .\\
 & & & & & & * & . & * & . & * & . & .\\
 & & & & & & * & . & * & . & * & . & .\\
0 & 0 & 0 & 1 & 0 & & * & . & * & . & * & . & .\\
0 & 1 & 0 & \binom{k}{1} & 0 & \cdots & * & . & * & . & * & . & .\\
1 & \binom{k}{1} & 0 & \binom{k}{2} & 0 & \cdots & * & . & * & . & * & . & .\\
0 & 0 & 1 & \binom{k}{3} & 0 & & * & . & * & . & * & . & .\\
0 & 0 & 0 & 0 & 1 & & * & . & * & . & * & . & .\\
 & & & & & & * & . & * & . & * & . & .\\
. & . & . & . & . & \cdots & * & . & * & . & * & . & .\\
. & . & . & . & . & & * & . & * & . & * & . & .\\
. & . & . & . & . & & * & . & * & . & * & . & .\\
 & & & & & & \binom{k}{k-7} & 0 & \binom{k}{k-6} & 0 & \binom{k}{k-5} & 0 & 0\\
 & & & & & & 0 & 1 & \binom{k}{k-5} & 0 & \binom{k}{k-4} & 0 & 0\\
 & & & & & & 0 & 0 & 0 & 1 & \binom{k}{k-3} & 0 & 0\\
 & & & & & & 0 & 0 & 0 & 0 & 0 & 1 & 0\\
0 & 0 & 0 & 0 & 0 & & 0 & 0 & 0 & 0 & 0 & 0 & 1
\end{pmatrix}$$

for k even.

$$A^{\beta}(S)=\begin{pmatrix}
0 & 0 & 0 & 0 & 0 & & & 0 & 0 & 0 & 0 & 1 & 0 & 0\\
 & & & & & & & 0 & 0 & 1 & 0 & \binom{k}{1} & 0 & 0\\
 & & & & & & & 1 & 0 & \binom{k}{1} & 0 & \binom{k}{2} & 0 & 0\\
 & & & & & & & \binom{k}{1} & 0 & \binom{k}{2} & 0 & \binom{k}{3} & 0 & 0\\
 & & & & & & & * & . & * & . & * & . & .\\
 & & & & & & \dots & * & . & * & . & * & . & .\\
 & & & & & & & * & . & * & . & * & . & .\\
 & & & & & & & * & . & * & . & * & . & .\\
0 & 0 & 0 & 0 & 1 & 0 & & * & . & * & . & * & . & .\\
0 & 0 & 1 & 0 & \binom{k}{1} & 0 & \dots & * & . & * & . & * & . & .\\
1 & 0 & \binom{k}{1} & 0 & \binom{k}{2} & 0 & \dots & * & . & * & . & * & . & .\\
0 & 1 & \binom{k}{2} & 0 & \binom{k}{3} & 0 & & * & . & * & . & * & . & .\\
0 & 0 & 0 & 1 & \binom{k}{4} & 0 & & * & . & * & . & * & . & .\\
0 & 0 & 0 & 0 & 0 & 1 & & * & . & * & . & * & . & .\\
. & . & . & . & . & & \dots & * & . & * & . & * & . & .\\
. & . & . & . & . & & & * & . & * & . & * & . & .\\
. & . & . & . & . & & & * & . & * & . & * & . & .\\
 & & & & & & & \binom{k}{k-7} & 0 & \binom{k}{k-6} & 0 & \binom{k}{k-5} & 0 & 0\\
 & & & & & & & 0 & 1 & \binom{k}{k-5} & 0 & \binom{k}{k-4} & 0 & 0\\
 & & & & & & & 0 & 0 & 0 & 1 & \binom{k}{k-3} & 0 & 0\\
 & & & & & & & 0 & 0 & 0 & 0 & 0 & 1 & 0\\
0 & 0 & 0 & 0 & 0 & & & 0 & 0 & 0 & 0 & 0 & 0 & 1
\end{pmatrix}$$

for k odd.

The "$*$" means $\binom{k}{j}$, and j increases by one when we move downwards row by row.

Proof: We need some steps.

1) For an upper triangular Stokes' matrix S (with entries s_{ij}) the entries of the matrix A^{β} which are different from zero are the following:

k **odd:**

$$A^{\beta}_{k-1,k-1}=A^{\beta}_{k,k}=1$$

For j even, $2+2i\le j\le k-2$, and $i=1,2,...,\frac{k}{2}-1$

$$A^{\beta}_{\frac{k}{2}-i,2i}=1$$

$$A^{\beta}_{\frac{k}{2}-i,j}=-s_{2i,j}+\sum_{n=1}^{\frac{j}{2}-i-1}(-1)^{n+1}F_{n,i,j}$$

$$F_{n,i,j}=\sum_{\alpha_1=1}^{\frac{j}{2}-i-n}\left(\sum_{\alpha_2=1}^{\frac{j}{2}-i-\alpha_1-n+1}\left(\sum_{\alpha_3=1}^{\frac{j}{2}-i-\alpha_1-\alpha_2-n+2}\left(\dots\sum_{\alpha_p=1}^{\frac{j}{2}-i-\sum_{r=1}^{p-1}\alpha_r-n+p-1}\left(\dots\sum_{\alpha_n=1}^{\frac{j}{2}-i-\sum_{r=1}^{n-1}\alpha_r-1}f(i,j,\alpha)\right)\right)\right)\right)$$

$$f(i,j,\alpha)=s_{2i,2i+2\alpha_1}\ s_{2i+2\alpha_1,2i+2\alpha_1+2\alpha_2}\ s_{2i+2\alpha_1+2\alpha_2,2i+2\alpha_1+2\alpha_2+2\alpha_3}\ \dots\ s_{2i+2\alpha_1+\dots+2\alpha_n,j}$$

For j even, $2i+2\le j\le k-2$ and $i=0,1,...,\frac{k}{2}-2$

$$A^{\beta}_{\frac{k}{2}+i,2i+1}=1$$

$$A_{\frac{k}{2}+i,j}=-s_{2i+1,j}+\sum_{n=1}^{\frac{j}{2}-i-1}(-1)^{n+1}G_{n,i,j}$$

$$G_{n,i,j}=\sum_{\alpha_1=1}^{\frac{j}{2}-i-n}\left(\sum_{\alpha_2=1}^{\frac{j}{2}-i-\alpha_1-n+1}\left(\sum_{\alpha_3=1}^{\frac{j}{2}-i-\alpha_1-\alpha_2-n+2}\left(\dots\sum_{\alpha_p=1}^{\frac{j}{2}-i-\sum_{r=1}^{p-1}\alpha_r-n+p-1}\left(\dots\sum_{\alpha_n=1}^{\frac{j}{2}-i-\sum_{r=1}^{n-1}\alpha_r-1}g(i,j,\alpha)\right)\right)\right)\right)$$

$$g(i,j,\alpha)=s_{2i+1,2i+2\alpha_1}\ s_{2i+2\alpha_1,2i+2\alpha_1+2\alpha_2}\ s_{2i+2\alpha_1+2\alpha_2,2i+2\alpha_1+2\alpha_2+2\alpha_3}\ \dots\ s_{2i+2\alpha_1+\dots+2\alpha_n,j}$$

More explicitely,

$$A^{\beta}(S)=\begin{pmatrix}
. & . & . & . & . & . & ...\\
. & . & . & . & . & . & ...\\
. & . & . & . & 0 & 0 & ...\\
. & . & 0 & 0 & 0 & 1 & ...\\
. & 0 & 0 & 1 & 0 & -s_{46} & ...\\
0 & 1 & 0 & -s_{24} & 0 & -s_{26}+s_{24}s_{46} & ...\\
1 & -s_{12} & 0 & -s_{14}+s_{12}s_{24} & 0 & -s_{16}+(s_{12}s_{26}+s_{14}s_{46})-s_{12}s_{24}s_{46} & ...\\
0 & 0 & 1 & -s_{34} & 0 & -s_{36}+s_{34}s_{46} & ...\\
. & . & 0 & 0 & 1 & -s_{56} & ...\\
. & . & . & . & 0 & 0 & ...\\
. & . & . & . & . & . & ...\\
. & . & . & . & . & . & ...
\end{pmatrix} \tag{25}$$

k **odd:**

$$A_{k-1,k-1}=A_{k,k}=1$$

For j odd, $2i+1\leq j\leq k-2$, $i=1,...,\frac{k-3}{2}$:

$$A_{\frac{k-1}{2},2i}=1$$

$$A_{\frac{k-1}{2}+i,j}=-s_{2i,j}+\sum_{n=1}^{\frac{j-1}{2}-i}(-1)^{n+1}H(n,i,j)$$

$$H(n,i,j)=\sum_{\alpha_1=0}^{\frac{j-1}{2}-i-n}\left(\sum_{\alpha_2=1}^{\frac{j-1}{2}-i-\alpha_1-n+1}\left(...\sum_{\alpha_p=1}^{\frac{j-1}{2}-i-\sum_{\alpha_r=1}^{p-1}\alpha_r-n+p-1}\left(...\sum_{\alpha_n=1}^{\frac{j-1}{2}-i-\sum_{\alpha_r=1}^{n-1}\alpha_r-1}h(i,j,\alpha)\right)\right)\right)$$

$$h(i,j,\alpha)=s_{2i,2i+1+2\alpha_1}\ s_{2i+1+2\alpha_1,2i+1+2\alpha_1+2\alpha_2}\ ....\ s_{2i+1+2\alpha_1+...+2\alpha_n,j}$$

For j odd, $2i+3\leq j\leq k-2$, $i=0,1,...,\frac{k-3}{2}$:

$$A_{\frac{k-1}{2}-i,2i+1}=1$$

$$A_{\frac{k-1}{2}-i,j}=-s_{2i+1,j}+\sum_{n=0}^{\frac{j-1}{2}-i}(-1)^{n+1}V(n,i,j)$$

$$V(n,i,j)=\sum_{\alpha_1=1}^{\frac{j-1}{2}-i-n}\left(\sum_{\alpha_2=1}^{\frac{j-1}{2}-i-\alpha_1-n+1}\left(...\sum_{\alpha_p=1}^{\frac{j-1}{2}-i-\sum_{r=1}^{p-1}\alpha_r-n+p-1}\left(...\sum_{\alpha_n=1}^{\frac{j-1}{2}-i-\sum_{r=1}^{n-1}\alpha_r-1}v(i,j,\alpha)\right)\right)\right)$$

$$v(i,j,\alpha)=s_{2i+1,2i+1+2\alpha_1}\ s_{2i+1+2\alpha_1,2i+1+2\alpha_1+2\alpha_2}\ ...\ s_{2i+1+2\alpha_1+...+2\alpha_n,j}$$

More explicitely:

$$A^{\beta}(S)=\begin{pmatrix}
. & . & . & . & . & . & . & ...\\
. & . & . & . & . & . & 0 & ...\\
. & . & . & . & 0 & 0 & 1 & ...\\
. & . & 0 & 0 & 1 & 0 & -s_{57} & ...\\
0 & 0 & 1 & 0 & -s_{35} & 0 & -s_{37}+s_{35}s_{57} & ...\\
1 & 0 & -s_{13} & 0 & -s_{15}+s_{13}s_{35} & 0 & -s_{17}+(s_{13}s_{37}+s_{15}s_{57})-s_{13}s_{35}s_{57} & ...\\
0 & 1 & -s_{23} & 0 & -s_{25}+s_{23}s_{35} & 0 & -s_{27}+(s_{23}s_{37}+s_{25}s_{57})-s_{23}s_{35}s_{57} & ...\\
. & 0 & 0 & 1 & -s_{45} & 0 & -s_{47}+s_{45}s_{57} & ...\\
. & . & . & 0 & 0 & 1 & -s_{67} & ...\\
. & . & . & . & . & 0 & 0 & ...\\
. & . & . & . & . & . & . & ...\\
. & . & . & . & . & . & . & ...
\end{pmatrix} \tag{26}$$

In order to prove the above expressions, we have to find each matrix $A^{\beta_{i,i+1}}$ corresponding to the elementary braid $\beta_{i,i+1}$ appearing in β. This means computing its entry $(i+1,i+1)$. This is a rather complicated problem, since the entry $(i+1,i+1)$ of a given $A^{\beta_{i,i+1}}$ is minus the entry $(i,i+1)$ of the Stokes' matrix resulting from the action of the elementary braids acting before $\beta_{i,i+1}$, which in general is a sum of products of the elements s_{ij} of the initial Stokes matrix S.

First we recall that $S \mapsto A^{\beta_{i,i+1}}\ S\ A^{\beta_{i,i+1}}$ has the following effect on the entries of S:

$$s_{n,i} \mapsto s_{n,i+1}, \qquad s_{n,i+1} \mapsto s_{n,i} - s_{i,i+1}s_{n,i+1}, \qquad n = 1,2,...,i-1$$

$$s_{i,i+1} \mapsto -s_{i,i+1}$$

$$s_{i,n} \mapsto s_{i+1,n}, \qquad s_{i+1,n} \mapsto s_{i,n} - s_{i,i+1}s_{i+1,n}, \qquad n = i+2,...,k$$

while all the other entries of S remain unchanged.

We start from $A^{\beta_{k-5,k-4}}$, whose non trivial entry is simply $-s_{k-5,k-4}$. Its action on S brings $s_{k-6,k-5}$ to $s_{k-6,k-4}$ (the reader may compute all the elements of $S^{\beta_{k-5,k-4}}$).

Then, the entry of $A^{\beta_{k-6,k-5}}$ is $-s_{k-6,k-4}$. Proceeding in this way, the reader may check that for the first sequence of braids $(\beta_{k-5,k-4}\beta_{k-6,k-5}\ ...\ \beta_{12})$ the entries $(i,i+1)$ of the matrices are:

$$-s_{i,k-4} \ \text{ for } A^{\beta_{i,i+1}}$$

Now, observe that

$$\begin{pmatrix}
1 &&&&&&&& \\
& \ddots &&&&&&& \\
&& 1 &&&&&& \\
&&& 0 & 1 &&&& \\
&&& 1 & x_1 &&&& \\
&&&&& 1 &&& \\
&&&&&& 1 && \\
&&&&&&& \ddots & \\
&&&&&&&& 1
\end{pmatrix}
\begin{pmatrix}
1 &&&&&&&& \\
& \ddots &&&&&&& \\
&& 1 &&&&&& \\
&&& 1 &&&&& \\
&&&& 0 & 1 &&& \\
&&&& 1 & x_2 &&& \\
&&&&&& 1 && \\
&&&&&&& \ddots & \\
&&&&&&&& 1
\end{pmatrix}
=
\begin{pmatrix}
1 &&&&&&&& \\
& \ddots &&&&&&& \\
&& 1 &&&&&& \\
&&& 0 & 0 & 1 &&& \\
&&& 1 & 0 & x_1 &&& \\
&&& 0 & 1 & x_2 &&& \\
&&&&&& 1 && \\
&&&&&&& \ddots & \\
&&&&&&&& 1
\end{pmatrix}$$

and recall that $A^{\beta_1\beta_2} = A^{\beta_2}A^{\beta_1}$. This implies for the first sequence of braids:

$$m_1 := A^{\beta_{k-5,k-4}\ \beta_{k-6,k-5}\ \cdots\ \beta_{12}}
\begin{pmatrix}
0 &&&&& 1 &&&& \\
1 & 0 &&&& * &&&& \\
& 1 & 0 &&& * &&&& \\
&& \ddots & \ddots && \vdots &&&& \\
&&& \ddots & 0 & * &&&& \\
&&&& 1 & * &&&& \\
&&&&&& 1 &&& \\
&&&&&&& 1 && \\
&&&&&&&& 1 & \\
&&&&&&&&& 1
\end{pmatrix}$$

the entries $*$ are exactly those of the $A_{i,i+1}$'s, namely $-s_{1,k-4}$, $-s_{2,k-4}$, ..., $-s_{k-5,k-4}$ from the top to the bottom of the $(k-4)^{th}$ column.

For the second sequence of braids $(\beta_{k-7,k-6}\ ...\ \beta_{34})$, the entries are

$$-s_{i-1,k-6} \ \text{ for } A^{\beta_{i,i+1}}$$

and, as above:

$$m_2 := A^{\beta_{k-6,k-5}\ \beta_{k-7,k-6}\ \cdots\ \beta_{23}} =
\begin{pmatrix}
1 &&&&&& 0 &&&&& \\
& 0 &&&&& 1 &&&&& \\
& 1 & 0 &&&& * &&&&& \\
&& 1 & 0 &&& * &&&&& \\
&&& \ddots & \ddots && \vdots &&&&& \\
&&&& \ddots & 0 & * &&&&& \\
&&&&& 1 & * &&&&& \\
&&&&&&& 1 &&&& \\
&&&&&&&& 1 &&& \\
&&&&&&&&& 1 && \\
&&&&&&&&&& 1 & \\
&&&&&&&&&&& 1
\end{pmatrix}$$

and the entries $*$ are those of the $A_{i,i+1}$'s (namely, $-s_{1,k-6}$, ..., $-s_{k-7,k-6}$ from the top to the bottom).

For the third sequence $(\beta_{k-6,k-5}\ \dots\ \beta_{23})$, they are

$$-s_{i-2,k-8} \ \text{ for } A^{\beta_{i,i+1}}$$

and:

$$m_3 := A^{\beta_{k-7,k-6}\ \cdots\ \beta_{34}} = \begin{pmatrix} 1 &&&&&&& 0 &&&&&& \\ & 1 &&&&&& 0 &&&&&& \\ && 0 &&&&& 1 &&&&&& \\ && 1 & 0 &&&& * &&&&&& \\ &&& 1 & 0 &&& * &&&&&& \\ &&&& \ddots & \ddots && \vdots &&&&&& \\ &&&&& \ddots & 0 & * &&&&&& \\ &&&&&& 1 & * &&&&&& \\ &&&&&&&& 1 &&&&& \\ &&&&&&&&& 1 &&&& \\ &&&&&&&&&& 1 &&& \\ &&&&&&&&&&& 1 && \\ &&&&&&&&&&&& 1 & \\ &&&&&&&&&&&&& 1 \end{pmatrix}$$

And so on. We reach the last but one "sequence", namely $\beta_{\frac{k}{2}-2,\frac{k}{2}-1}$ for k even, or $(\beta_{\frac{k-3}{2},\frac{k-1}{2}}\beta_{\frac{k-5}{2},\frac{k-3}{2}})$ for k odd. The entries are $-s_{12}$, or $-s_{23}$,$-s_{13}$ respectively. Then

$$m_{\frac{k}{2}-2} := A^{\beta_{\frac{k}{2}-2,\frac{k}{2}-1}} = \begin{pmatrix} 1 &&&&&&& \\ & \ddots &&&&&& \\ && 1 &&&&& \\ &&& 0 & 1 &&& \\ &&& 1 & -s_{12} &&& \\ &&&&& 1 && \\ &&&&&& \ddots & \\ &&&&&&& 1 \end{pmatrix}$$

or

$$m_{\frac{k-5}{2}} := A^{\beta_{\frac{k-3}{2},\frac{k-1}{2}}\beta_{\frac{k-5}{2},\frac{k-3}{2}}} = \begin{pmatrix} 1 &&&&&&&& \\ & \ddots &&&&&&& \\ && 1 &&&&&& \\ &&& 0 & 0 & 1 &&& \\ &&& 1 & 0 & -s_{13} &&& \\ &&&& 1 & -s_{23} &&& \\ &&&&&& 1 && \\ &&&&&&& \ddots & \\ &&&&&&&& 1 \end{pmatrix}$$

The entries for the last braid are more complicated, because the entries on the first upper sub-diagonal of the Stokes' matrix have been shuffled by the preceeding braids. We give the result ($A_{i,i+1}$ stends for $A^{\beta_{i,i+1}}$

$$A_{k-3,k-2}: \quad -s_{k-3,k-2}$$

$$A_{k-4,k-3}: \quad -s_{k-5,k-2}+s_{k-5,k-4}s_{k-4,k-2}$$

$$A_{k-5,k-4}: \quad -s_{k-7,k-2}+(s_{k-7,k-4}s_{k-4,k-2}+s_{k-7,k-6}s_{k-6,k-2})-s_{k-7,k-6}s_{k-6,k-4}s_{k-4,k-2}$$

$$\vdots$$

$$A_{\frac{k}{2}-1,\frac{k}{2}}: \quad -s_{1,k-2}+(s_{12}s_{2,k-2}+s_{14}s_{4,k-2}+\dots+s_{1,k-4}s_{k-4,k-2})\ +\ \dots\ +\ (-1)^{\frac{k}{2}-1}s_{12}s_{24}s_{48}\dots s_{k-4,k-2}$$

$$A_{\frac{k}{2}-2,\frac{k}{2}-1}: \quad -s_{2,k-2}+(s_{24}s_{4,k-2}+s_{26}s_{4,k-2}+\dots+s_{2,k-4}s_{k-4,k-2})\ +\ \dots\ +\ (-1)^{\frac{k}{2}-2}s_{24}s_{48}\dots s_{k-4,k-2}$$

$$\vdots$$

$$A_{34}: \quad -s_{k-8,k-2}+(s_{k-8,k-4}s_{k-4,k-2}+s_{k-8,k-6}s_{k-6,k-2})-s_{k-8,k-6}s_{k-6,k-4}s_{k-4,k-2}$$

$$A_{23}: \quad -s_{k-6,k-2}+s_{k-6,k-4}s_{k-4,k-2}$$

$$A_{12}: \quad -s_{k-4,k-2}$$

For k odd

$$A_{k-3,k-2}: \quad -s_{k-3,k-2}$$

$$A_{k-4,k-3}: \quad -s_{k-5,k-2}+s_{k-5,k-4}s_{k-4,k-2}$$

$$A_{k-5,k-4}: \quad -s_{k-7,k-2}+(s_{k-7,k-4}s_{k-4,k-2}+s_{k-7,k-6}s_{k-6,k-2})-s_{k-7,k-6}s_{k-6,k-4}s_{k-4,k-2}$$

$$\vdots$$

$$A_{\frac{k-1}{2},\frac{k+1}{2}}: \quad -s_{2,k-2}+(s_{23}s_{3,k-2}+s_{25}s_{5,k-2}+...+s_{2,k-4}s_{k-4,k-2})\ +\ ...\ +\ (-1)^{\frac{k-3}{2}}s_{23}s_{35}s_{57}...s_{k-4,k-2}$$

$$A_{\frac{k-3}{2},\frac{k-1}{2}}: \quad -s_{1,k-2}+(s_{13}s_{3,k-2}+s_{15}s_{5,k-2}+...+s_{1,k-4}s_{k-4,k-2})\ +\ ...\ +\ (-1)^{\frac{k-3}{2}}s_{13}s_{35}s_{57}...s_{k-4,k-2}$$

$$\vdots$$

$$A_{34}: \quad -s_{k-8,k-2}+(s_{k-8,k-4}s_{k-4,k-2}+s_{k-8,k-6}s_{k-6,k-2})-s_{k-8,k-6}s_{k-6,k-4}s_{k-4,k-2}$$

$$A_{23}: \quad -s_{k-6,k-2}+s_{k-6,k-4}s_{k-4,k-2}$$

$$A_{12}: \quad -s_{k-4,k-2}$$

and:

$$A^{\beta_{k-3,k-2}\ \beta_{k-4,k-3}\ \cdots\ \beta_{12}}=\begin{pmatrix} 0 & & & & & 1 & & \\ 1 & 0 & & & & * & & \\ & 1 & 0 & & & * & & \\ & & \ddots & \ddots & & \vdots & & \\ & & & \ddots & 0 & * & & \\ & & & & 1 & * & & \\ & & & & & & 1 & 0 \\ & & & & & & 0 & 1 \end{pmatrix}$$

which we call $m_{\frac{k}{2}-1}$ for k even, $m_{\frac{k-3}{2}}$ for k odd.

Then, for k even

$$A^{\beta}=m_{\frac{k}{2}-1}m_{\frac{k}{2}-2}\ ...\ m_3m_2m_1$$

and for k odd

$$A^{\beta}=m_{\frac{k-3}{2}}m_{\frac{k-5}{2}-2}\ ...\ m_3m_2m_1$$

Doing a careful computation, we obtain (25) and (26).

2) The second step consists of expressing A^{β} in terms of the entries of the Stokes' factors of S, which are simply binomial coefficients. First we prove the following

Lemma 9: *Given an upper triangular $k\times k$ matrix S, with entries $s_{ii}=1$, we can uniquely determine numbers a_{ij} such that, for k, i, j all even or all odd:*

$$s_{ij}=a_{ij}\ +\ (a_{i,i+2}a_{i+2,j}+a_{i,i+4}a_{i+4,j}+\ ...\ +a_{i,j-2}a_{j-2,j})\ +$$

$$+\ (a_{i,i+2}a_{i+2,,i+4}a_{i+4,j}+a_{i,i+2}a_{i+2,i+6}a_{i+6,j}+\ ...\ +a_{i,j-4}a_{j-4,j-2}a_{j-2,j})\ +....\ +\ a_{i,i+2}a_{i+2,i+4}a_{i+4,i+6}\ ...\ a_{j-2,j}$$

If k is even, but i is odd, just replace in the formula $i+2$ with $i+1$, $i+4$ with $i+3$, ect. If k is even, but j is odd, just replace $j-2$ with $j-1$, $j-4$ with $j-3$, ect.

If k is odd, but i is even, or j is even, just do the same replacements as above. More explicitely:

$$S:=\begin{pmatrix} 1 & a_{12} & a_{13}+a_{12}a_{23} & a_{14}+a_{12}a_{24} & \begin{matrix} a_{15}+(a_{12}a_{25}+a_{14}a_{45}) \\ +a_{12}a_{24}a_{45} \end{matrix} & \begin{matrix} a_{16}+(a_{12}a_{26}+a_{14}a_{46}) \\ +a_{12}a_{24}a_{46} \end{matrix} & \begin{matrix} a_{17}+(a_{12}a_{27}+a_{14}a_{47}+a_{16}a_{67}) \\ +(a_{12}a_{24}a_{47}+a_{12}a_{26}a_{67}+a_{14}a_{46}a_{67}) \\ +a_{12}a_{24}a_{46}a_{67} \end{matrix} & .... \\ & 1 & a_{23} & a_{24} & a_{25}+a_{24}a_{45} & a_{26}+a_{24}a_{46} & a_{27}+(a_{24}a_{47}+a_{26}a_{67})+a_{24}a_{46}a_{67} & .... \\ & & 1 & a_{34} & a_{35}+a_{34}a_{45} & a_{36}+a_{34}a_{46} & a_{37}+(a_{34}a_{47}+a_{36}a_{67})+a_{34}a_{46}a_{67} & .... \\ & & & 1 & a_{45} & a_{46} & a_{47}a_{46}a_{67} & .... \\ & & & & 1 & a_{56} & a_{57}+a_{56}a_{57} & .... \\ & & & & & 1 & a_{67} & .... \\ & & & & & & 1 & .... \\ & & & & & & & .... \end{pmatrix}$$

for k even.

$$
S := \begin{pmatrix}
1 & a_{12} & a_{13} & a_{14} + a_{13}a_{34} & a_{15} + a_{13}a_{35} & a_{16} + (a_{13}a_{36} + a_{15}a_{56}) + a_{13}a_{35}a_{56} & a_{17} + (a_{13}a_{37} + a_{15}a_{57}) + a_{13}a_{35}a_{57} & \dots \\
 & 1 & a_{23} & a_{24} + a_{23}a_{34} & a_{25} + a_{23}a_{35} & a_{26} + (a_{23}a_{36} + a_{25}a_{56}) + a_{23}a_{35}a_{56} & a_{27} + (a_{23}a_{37} + a_{25}a_{57}) + a_{23}a_{35}a_{57} & \dots \\
 & & 1 & a_{34} & a_{35} & a_{36} + a_{35}a_{56} & a_{37} + a_{35}a_{57} & \dots \\
 & & & 1 & a_{45} & a_{46} + a_{45}a_{56} & a_{47} + a_{45}a_{57} & \dots \\
 & & & & 1 & a_{56} & a_{57} & \dots \\
 & & & & & 1 & a_{67} & \dots \\
 & & & & & & 1 & \dots \\
 & & & & & & & \dots
\end{pmatrix}
$$

for k odd.

Proof: We have to solve a non linear system $F_{ij}(a) = s_{ij}$.

The sum of the differences between the indices of the factors a in s_{ij} is equal to the difference of the indices of s_{ij}, namely $j - i$.

¿From this it follows that the trems non-linear in the a's occur on sub-diagonals which lie above all the sub-diagonals containing the factors of the non linear terms.

Then, the system $F_{ij}(a) = s_{ij}$ is uniquely solvable, starting from the first sub-diagonal and successively determining all the a_{rs} going up diagonal by diagonal.

□

Corollary 3: *With the above factorization, the matrix A^{β} becomes:*

$$
A^{\beta}(S) = \begin{pmatrix}
. & . & . & . & . & . & . & . & \dots \\
. & . & . & . & . & . & . & 0 & \dots \\
. & . & . & . & . & 0 & 0 & 1 & \dots \\
. & . & . & 0 & 0 & 1 & 0 & -a_{68} & \dots \\
. & 0 & 0 & 1 & 0 & -a_{46} & 0 & -a_{48} & \dots \\
0 & 1 & 0 & -a_{24} & 0 & -a_{26} & 0 & -a_{28} & \dots \\
1 & -a_{12} & 0 & -a_{14} & 0 & -a_{16} & 0 & -a_{18} & \dots \\
0 & 0 & 1 & -a_{34} & 0 & -a_{36} & 0 & -a_{38} & \dots \\
. & . & 0 & 0 & 1 & -a_{56} & 0 & -a_{58} & \dots \\
. & . & . & . & 0 & 0 & 1 & -a_{78} & \dots \\
. & . & . & . & . & . & 0 & 0 & \dots \\
. & . & . & . & . & . & . & . & \dots
\end{pmatrix} \tag{27}
$$

for k even; and

$$
A^{\beta}(S) = \begin{pmatrix}
. & . & . & . & . & . & . & \dots \\
. & . & . & . & . & . & 0 & \dots \\
. & . & . & . & 0 & 0 & 1 & \dots \\
. & . & 0 & 0 & 1 & 0 & -a_{57} & \dots \\
0 & 0 & 1 & 0 & -a_{35} & 0 & -a_{37} & \dots \\
1 & 0 & -a_{13} & 0 & -a_{15} & 0 & -a_{17} & \dots \\
0 & 1 & -a_{23} & 0 & -a_{25} & 0 & -a_{27} & \dots \\
. & 0 & 0 & 1 & -a_{45} & 0 & -a_{47} & \dots \\
. & . & . & 0 & 0 & 1 & -a_{67} & \dots \\
. & . & . & . & . & 0 & 0 & \dots \\
. & . & . & . & . & . & . & \dots \\
. & . & . & . & . & . & . & \dots
\end{pmatrix} \tag{28}
$$

for k odd. In other words, all the non linear terms in formulae (25) and (26) drop.

Proof: Just substitute the factorization of S in (25), (26).

□

In section 5 we computed the Stokes' factors for S. If we sum all the factors appearing in formula

(20) we get a matrix of the form:

$$M := K_{1k} + K_{1,k-1} + K_{k,k-1} + ... + K_{k3} + K_{k2} = \begin{pmatrix} k & & & & & & 0 \\ * & k & & & & 0 & * \\ * & * & k & & & * & * \\ \vdots & \vdots & \vdots & \ddots & \iddots & \vdots & \vdots \\ * & * & * & \iddots & \ddots & \vdots & \vdots \\ * & * & 0 & & & k & * \\ * & 0 & & & & & k \end{pmatrix}$$

The $*$ are the binomial coefficient appearing in the factors. If we know M, we can determine all the entries of the single Stokes' factors, because if the entry (i,j) is not zero for one factor, then it is zero for all the other factors. Now, we rename the entries of the factors according to the following rule:

$$M := \begin{pmatrix} k & & & & & & & & & & & 0 \\ a_{k-2,k} & k & & & & & & & & & 0 & a_{k-2,k-1} \\ a_{k-4,k} & a_{k-4,k-2} & k & & & & & & & & a_{k-4,k-3} & a_{k-4,k-1} \\ \vdots & \vdots & \vdots & \ddots & & & & & & \iddots & \vdots & \vdots \\ a_{4k} & a_{4,k-2} & ... & a_{46} & k & 0 & 0 & 0 & a_{45} & a_{47} & ... & a_{4,k-1} \\ a_{2,k} & a_{2,k-2} & ... & a_{26} & a_{24} & k & 0 & a_{23} & a_{25} & a_{27} & ... & a_{2,k-1} \\ a_{1k} & a_{1,k-2} & ... & a_{16} & a_{14} & a_{12} & k & a_{13} & a_{15} & a_{17} & ... & a_{1.k-1} \\ a_{3k} & a_{3,k-2} & ... & a_{36} & a_{34} & 0 & 0 & k & a_{35} & a_{37} & ... & a_{3,k-1} \\ a_{5k} & a_{5,k-2} & ... & a_{56} & 0 & 0 & 0 & 0 & k & a_{57} & ... & a_{5,k-1} \\ \vdots & \vdots & \iddots & & & & & & & \ddots & ... & ... \\ a_{k-3,k} & a_{k-3,k-2} & & & & & & & & & k & a_{k-3,k-1} \\ a_{k-1,k} & & & & & & & & & & & 1 \end{pmatrix}$$

for k even.

$$M = \begin{pmatrix} k & & & & & & & & & & & & 0 \\ a_{k-2,k} & k & & & & & & & & & & & a_{k-2,k-1} \\ a_{k-4,k} & a_{k-4,k-2} & k & & & & & & & & & a_{k-4,k-3} & a_{k-4,k-1} \\ \vdots & \vdots & \vdots & \ddots & & & & & & & \iddots & \vdots & \vdots \\ a_{5k} & a_{5,k-2} & a_{5,k-4} & ... & k & 0 & 0 & 0 & 0 & a_{56} & ... & a_{5,k-3} & a_{5,k-1} \\ a_{3k} & a_{3,k-2} & a_{3,k-4} & ... & a_{35} & k & 0 & 0 & a_{34} & a_{36} & ... & a_{3,k-3} & a_{3,k-1} \\ a_{1k} & a_{1,k-2} & a_{1,k-4} & ... & a_{15} & a_{13} & k & a_{12} & a_{14} & a_{16} & ... & a_{1,k-3} & a_{1,k-1} \\ a_{2k} & a_{2,k-2} & a_{2,k-4} & ... & a_{25} & a_{23} & 0 & k & a_{24} & a_{26} & ... & a_{2,k-3} & a_{2,k-1} \\ a_{4k} & a_{4,k-2} & a_{4,k-4} & ... & a_{45} & 0 & 0 & 0 & k & a_{46} & ... & a_{4,k-3} & a_{4,k-1} \\ \vdots & \vdots & & \iddots & & & & & & & \ddots & \vdots & \vdots \\ a_{k-3,k} & a_{k-3,k-2} & & & & & & & & & & k & a_{k-3,k-1} \\ a_{k-1,k} & & & & & & & & & & & & k \end{pmatrix}$$

for k odd. The strange labelling is simply the one such that

$$PMP^{-1} = \begin{pmatrix} k & a_{12} & a_{13} & a_{14} & ... \\ & k & a_{23} & a_{24} & ... \\ & & k & a_{34} & ... \\ & & & \ddots & \vdots \\ & & & & k \end{pmatrix}$$

where the matrix of permutation is

$$P = \begin{pmatrix} & & & & & & 1 & 0 & & & & & \\ & & & & & 1 & 0 & 0 & & & & & \\ & & & & 0 & 0 & 0 & 1 & 0 & & & & \\ & & & & 1 & 0 & 0 & 0 & 0 & & & & \\ & & & 0 & 0 & 0 & 0 & 0 & 1 & 0 & & & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ & 1 & & & & & & & & & & 0 & \\ 0 & 0 & & & & & & & & & & 0 & 1 \\ 1 & 0 & & & & & & & & & & 0 & 0 \end{pmatrix}$$

for k even (the 1 on the first row is on the $\frac{k}{2}+1$-th column) ; and

$$P = \left(\begin{array}{ccccccccccccc}
 & & & & & 0 & 1 & & & & & & \\
 & & & & & 0 & 0 & 1 & & & & & \\
 & & & & 0 & 1 & 0 & 0 & 0 & & & & \\
 & & & & 0 & 0 & 0 & 0 & 1 & & & & \\
 & & & 0 & 1 & 0 & 0 & 0 & 0 & 0 & & & \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
0 & 1 & & & & & & & & & & 0 & \\
0 & 0 & & & & & & & & & & 0 & 1 \\
1 & 0 & & & & & & & & & & 0 & 0
\end{array}\right)$$

for k odd (the 1 on the first row is on the $\frac{k+1}{2}$-th column).

With this choice of the labelling, the product $S_{upper} := P\,(K_{1k}K_{1,k-1}K_{k,k-1}\ ...\ K_{k3}K_{k2}\ )\,P^{-1}$ is precisely factorized as in lemma 9.

Then we can write the entries of A^{β} (formulae (27) (28)) from the entries of the Stokes factors (which are binomial coefficient).

The final result is precisely the claim of the proposition.

□

We are ready to prove the main result:

Theorem 2: *Consider the Stokes matrix* $S = T_F^{\frac{k}{2}}\ T^{-\frac{k}{2}}$ *(k even) or* $S = T_F^{\frac{k-1}{2}} K_{k2} T^{-\frac{k-1}{2}}$ *(k odd) for the quantum cohomology of* $\mathbf{CP}^{k-1}$ *and set it in the upper triangular form* $S_{upper} = P\ S\ P^{-1}$ *by the permutation* P*. Then, there exists a braid* β *(Lemma 8), represented by a matrix* A^{β} *(Proposition), which sets* S_{upper} *in the form (24). The last column is negative, but conjugation by* $\mathcal{I} = \mathrm{diag}(1,1,...,1,-1)$ *makes it positive. We reach the canonical form:*

$$s_{ij} = \binom{k}{j-i}, \qquad i<j$$

Another conjugation by $\mathrm{diag}(-1,1,-1,1,-1,...)$ *brings the matrix in the equivalent canonical form*

$$s_{ij} = (-1)^{j-i}\binom{k}{j-i}, \qquad i<j$$

Finally, by the action of the braid group, the last matrix can be put in the form

$$s_{ij} = \binom{k-1+j-i}{j-i}, \qquad i<j$$

In all the above matrices

$$s_{ii} = 1, \qquad s_{ij} = 0 \qquad i>j$$

Proof: First, we want to explain which is the braid which brings the upper triangular matrix with entries $s_{ij} = (-1)^{j-i}\binom{k}{j-i}$ in the matrix $s_{ij} = \binom{k-1+j-i}{j-i}$. We make use of the following known result [25]:

Consider the upper triangular Stokes' matrix S*, the braid* $\beta = \beta_{12}(\beta_{23}\beta_{12})(\beta_{34}\beta_{23}\beta_{12})\ ...\ \ (\beta_{n-1,n}\beta_{n-2,n-1}...\beta_{23}\beta_{12})$ *and the permutation*

$$P = \begin{pmatrix} & & & 1 \\ & & 1 & \\ & \cdot^{\cdot^{\cdot}} & & \\ 1 & & & \end{pmatrix}$$

Then, the relation

$$\left[S^{-1}\right]^{\beta} = PS^{T}P \tag{29}$$

holds.

Observe that for the matrix S, whose upper triangular part has entries $s_{ij} = \begin{pmatrix} k-1+j-i \\ j-i \end{pmatrix}$, we have $PS^TP \equiv S$. Moreover, S^{-1} is upper triangular with entries $s_{ij} = (-1)^{j-i}\begin{pmatrix} k \\ j-i \end{pmatrix}$. This proves that S and S^{-1} are equivalent w.r.t the action of the braid group.

Let us now prove the theorem staring from k even. We have to prove that $A^\beta\ P\ T_F^{\frac{k}{2}}\ T^{-\frac{k}{2}}\ P^{-1}[A^\beta]^T$ is in "canonical form" (24). The proof "reduces" to the computation of products of matrices explicitely given. We do the products in an shrewd way. First we rewrite

$$S^\beta = A^\beta\ \left(P\ T_F\ P^{-1}\right)^{\frac{k}{2}}\ \left(P\ T^{-1}\ P^{-1}\right)^{\frac{k}{2}}\ [A^\beta]^T$$

and we compute:

$$P\ T_F\ P^{-1} = \begin{pmatrix}
 & & & & & & 1 & 0 & & & & & \\
 & & & & & 1 & 0 & 0 & & & & & \\
 & & & & 0 & 0 & 0 & 1 & 0 & & & & \\
 & & & & 1 & 0 & 0 & 0 & 0 & & & & \\
 & & & 0 & 0 & 0 & 0 & 0 & 1 & 0 & & & \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
 & 1 & & & & & & & & & & 0 & \\
0 & 0 & & & & & & & & & & 0 & 1 \\
1 & 0 & & & & & & & & & & 0 & 0
\end{pmatrix}
\begin{pmatrix}
0 & & & & \cdots & 1 \\
-1 & 0 & & & & \\
 & -1 & 0 & & & \\
 & & -1 & & & \\
 & & & \ddots & \ddots & \vdots \\
 & & & & -1 & 0
\end{pmatrix}
\begin{pmatrix}
 & & & & & & 0 & 1 \\
 & & & & & 1 & 0 & 0 \\
 & & & & \vdots & & & \\
 & & 0 & 1 & & & & \\
 & 1 & 0 & 0 & & & & \\
1 & 0 & 0 & 0 & & & & \\
0 & 0 & 1 & 0 & & & & \\
 & & 0 & 0 & & & & \\
 & & & & \vdots & & & \\
 & & & & & 0 & 0 & 0 \\
 & & & & & & 1 & 0
\end{pmatrix}$$

(in P^{-1} the 1 on the first column is on the $\frac{k}{2}+1$-th row)

$$= \begin{pmatrix}
0 & -1 & & & & & & & & & & \\
0 & 0 & 0 & -1 & & & & & & & & \\
-1 & 0 & 0 & 0 & & & & & & & & \\
 & 0 & 0 & 0 & 0 & -1 & & & & & & \\
 & & -1 & 0 & 0 & 0 & & & & & & \\
 & & & & & & \vdots & \vdots & & & & \\
 & & & & & & & & 0 & 0 & 0 & -1 \\
 & & & & & & & & -1 & 0 & 0 & 0 \\
 & & & & & & & & & & 1 & 0
\end{pmatrix}$$

Thus

$$\left(P\ T_F\ P^{-1}\right)^{\frac{k}{2}} = \begin{pmatrix}
 & & & & & & & -1 \\
 & & & & & & 1 & \\
 & & & & & -1 & & \\
 & & & & \iddots & & & \\
 & & & -1 & & & & \\
 & & 1 & & & & & \\
 & -1 & & & & & & \\
1 & & & & & & &
\end{pmatrix}$$

Then, using the expression of A^β from the proposition:

$$F := A^\beta\ \left(P\ T_F\ P^{-1}\right)^{\frac{k}{2}} = \begin{pmatrix}
0 & 0 & 1 & & & & & & & & & \\
 & 0 & \binom{k}{1} & 0 & 1 & & & & & & & \\
 & & \binom{k}{2} & 0 & \binom{k}{1} & & & & & & & \\
 & & \binom{k}{3} & 0 & \binom{k}{2} & 0 & 1 & & & & & \\
 & & \vdots & & \vdots & & \vdots & & & & & \\
 & & & & & & & \cdots & 1 & & & \\
 & & * & & * & & * & \cdots & \binom{k}{1} & 0 & 1 & \\
 & & * & & * & & * & \cdots & \binom{k}{2} & 0 & \binom{k}{1} & -1 \\
 & & * & & * & & * & \cdots & \binom{k}{3} & -1 & 0 & \\
 & & \vdots & & \vdots & & \vdots & & & & & \\
 & & \binom{k}{k-4} & 0 & \binom{k}{k-5} & & & & & & & \\
 & & \binom{k}{k-3} & -1 & 0 & & & & & & & \\
0 & -1 & 0 & & & & & & & & & \\
-1 & 0 & & & & & & & & & &
\end{pmatrix}$$

(the -1 on the last column is on the $\frac{k}{2}$-th row).

Using the explicit expressions for K_{k3} and K_{k2} of section 5 we compute

$$T^{-1} = T_F^{-1}\ K_{k3}\ K_{k2} = \begin{pmatrix} 0 & -1 & & & & \\ & 0 & -1 & & & \\ & & \ddots & \ddots & & \\ & & & 0 & -1 & \\ & & & & 0 & -1 \\ 1 & & & & & 0 \end{pmatrix} K_{k3}\ K_{k2} =$$

$$= \begin{pmatrix} \binom{k}{1} & -1 & & & & & & & & & & & -\binom{k}{2} \\ & 0 & -1 & & & & & & & & & -\binom{k}{4} & -\binom{k}{3} \\ & & 0 & -1 & & & & & & & -\binom{k}{6} & -\binom{k}{5} & 0 \\ & & & \ddots & \ddots & & & & & \iddots & \iddots & & \\ & & & & 0 & -1 & 0 & -\binom{k}{k-2} & -\binom{k}{k-3} & & & & \\ & & & & & 0 & -1 & -\binom{k}{k-1} & & & & & \\ & & & & & & 0 & -1 & & & & & \\ & & & & & & & 0 & -1 & & & & \\ & & & & & & & & \ddots & \ddots & & & \\ & & & & & & & & & & & & \\ & & & & & & & & & & & 0 & -1 \\ & & & & & & & & & & & & 0 \end{pmatrix}$$

Then

$$P\ T^{-1}\ P^{-1} = \begin{pmatrix} 0 & 0 & -1 & & & & & & & & & & \\ -1 & 0 & -\binom{k}{k-1} & & & & & & & & & & \\ 0 & 0 & 0 & 0 & -1 & & & & & & & & \\ 0 & -1 & -\binom{k}{k-2} & 0 & -\binom{k}{k-3} & 0 & & & & & & & \\ & 0 & 0 & 0 & 0 & 0 & -1 & & & & & & \\ & & & -1 & -\binom{k}{k-4} & 0 & -\binom{k}{k-5} & & & & & & \\ & & & & & & & \ddots & & & & & \\ & & & & & & & & 0 & 0 & 0 & -1 & 0 \\ & & & & & & & & -1 & -\binom{k}{4} & 0 & -\binom{k}{3} & 0 \\ & & & & & & & & & 0 & 0 & 0 & 1 \\ & & & & & & & & & 0 & -1 & -\binom{k}{2} & \binom{k}{1} \end{pmatrix}$$

After this we computed $F\ \left[P\ T^{-1}\ P^{-1}\right]$, $F\ \left[P\ T^{-1}\ P^{-1}\right]\ \left[P\ T^{-1}\ P^{-1}\right]$, $F\ \left[P\ T^{-1}\ P^{-1}\right]^3$, ..., $F\ \left[P\ T^{-1}\ P^{-1}\right]^{\frac{k}{2}}$. We omit the intermediate computations and we give the final result:

$$F_1 := F\ \left(P\ T^{-1}\ P^{-1}\right)^{\frac{k}{2}} = \begin{pmatrix} & & & & & & & & & & 1 & \binom{k}{2} & -\binom{k}{1} \\ & & & & & & & & & \binom{k}{4} & 0 & \binom{k}{3} & -\binom{k}{2} \\ & & & & & & & & & \binom{k}{5} & 0 & \binom{k}{4} & -\binom{k}{3} \\ & & & & & & & \dots & & \vdots & \vdots & \vdots & \vdots \\ & & & & & 1 & \binom{k}{k-6} & \dots & \dots & * & 0 & * & * \\ & & & 1 & \binom{k}{k-4} & 0 & \binom{k}{k-5} & \dots & \dots & * & 0 & * & * \\ & 1 & \binom{k}{k-2} & 0 & \binom{k}{k-3} & 0 & \binom{k}{k-4} & \dots & \dots & * & 0 & * & * \\ 1 & 0 & \binom{k}{k-1} & 0 & \binom{k}{k-2} & 0 & \binom{k}{k-3} & \dots & \dots & * & 0 & * & * \\ & & 1 & 0 & \binom{k}{k-1} & 0 & \binom{k}{k-2} & \dots & \dots & * & 0 & * & * \\ & & & & 1 & 0 & \binom{k}{k-1} & \dots & \dots & * & 0 & * & * \\ & & & & & & 1 & \dots & \dots & * & 0 & * & * \\ & & & & & & & \dots & \dots & \vdots & \vdots & \vdots & \vdots \\ & & & & & & & & & \binom{k}{k-1} & 0 & \binom{k}{k-2} & -\binom{k}{k-4} \\ & & & & & & & & & & 1 & \binom{k}{k-1} & -\binom{k}{k-2} \\ & & & & & & & & & & & & -\binom{k}{k-1} \\ & & & & & & & & & & & & 1 \end{pmatrix}$$

Now, multiplying $F_1\ \left[A^{\beta}\right]^T$, we obtain precisely the “canonical form” (24).

For k odd, we did a similar computation. We omit the detail, but we indicate the order of multiplications which yielded the most simple expressions to multiply step by step. Our aim is to compute $A^\beta\ P\ T_F^{\frac{k-1}{2}}\ K_{k2}\ (T_F^{-1}\ K_{k3}\ K_{k2})^{\frac{k-1}{2}}\ P^{-1}\ [A^\beta]^T$. First, we computed PT_FP^{-1}, then $(PT_FP^{-1})^{\frac{k-1}{2}}$, then $F := A^\beta\ (PT_FP^{-1})^{\frac{k-1}{2}}$. After this, we computed $PK_{k2}P^{-1}$, and $F_1 := F\ PK_{k2}P^{-1}$. Finally, we calculated $m := P\ T_F^{-1}\ K_{k3}\ K_{k2}\ P^{-1}$ and $F_1\ m$, $F_1\ m\ m$, ..., $F_2 := F_1\ m^{\frac{k-1}{2}}$. The matrix $F_2\ [A^\beta]^T$ proved to be in "canonical form".

$\square$

7 Canonical form of S^{-1}

The matrix S^{-1} such that $Y_R(z) = Y_L(z)\ S^{-1}$ can be put in the same canonical form of S, as a consequence of the relation (29). The only remarks we want to add concern the braid which brings S^{-1} to the canonical form, because it arranges the lines L_j in a "beautiful" shape.

Lemma $8'$**:** *Let the points u_j be in lexicographical order w.r.t the admissible line l. Let us denote $\sigma_{i,i+1} := \beta_{i,i+1}^{-1}$. Then, the following braid arranges the poins in cyclic clockwise order, u_1 being the first point in Π_L for k even, or the last in Π_R for k odd (w.r.t the clockwise order) (see figure 6):*

$$\beta' := \Big[\ (\sigma_{34}\sigma_{56}\sigma_{78}\ ...\ \sigma_{k-3,k-2}\sigma_{k-1,k})\ (\sigma_{45}\sigma_{67}\sigma_{89}\ ...\ \sigma_{k-4,k-3}\sigma_{k-2,k-1})\ ...\ (\sigma_{\frac{k}{2},\frac{k}{2}+1}\sigma_{\frac{k}{2}+2,\frac{k}{2}+3})\ \sigma_{\frac{k}{2}+1,\frac{k}{2}+2}\Big]$$

$$\Big[\ (\sigma_{\frac{k}{2}+2,\frac{k}{2}+3}\sigma_{\frac{k}{2}+3,\frac{k}{2}+4}\ \cdots\ \sigma_{k-2,k-1}\sigma_{k-1,k})\ (\sigma_{\frac{k}{2}+2,\frac{k}{2}+3}\sigma_{\frac{k}{2}+3,\frac{k}{2}+4}\ \cdots\ \sigma_{k-2,k-1})\ ....\ (\sigma_{\frac{k}{2}+2,\frac{k}{2}+3}\sigma_{\frac{k}{2}+3,\frac{k}{2}+4})\ \sigma_{\frac{k}{2}+2,\frac{k}{2}+3}\Big]$$

for k even, and

$$\beta' := \beta_{12}\ \Big[\ (\sigma_{45}\sigma_{67}\sigma_{89}\ ...\ \sigma_{k-3,k-2}\sigma_{k-1,k})\ (\sigma_{56}\sigma_{78}\ ...\ \sigma_{k-4,k-3}\sigma_{k-2,k-1})\ ....\ (\sigma_{\frac{k+1}{2},\frac{k+3}{2}}\sigma_{\frac{k+5}{2},\frac{k+7}{2}})\ \sigma_{\frac{k+3}{2},\frac{k+5}{2}}\Big]$$

$$\Big[\ (\sigma_{\frac{k+5}{2},\frac{k+7}{2}}\sigma_{\frac{k+7}{2},\frac{k+9}{2}}\ ...\ \sigma_{k-2,k-1}\sigma_{k-1,k})\ (\sigma_{\frac{k+5}{2},\frac{k+7}{2}}\sigma_{\frac{k+7}{2},\frac{k+9}{2}}\ ...\ \sigma_{k-2,k-1})\ ....\ (\sigma_{\frac{k+5}{2},\frac{k+7}{2}}\sigma_{\frac{k+7}{2},\frac{k+9}{2}})\ \sigma_{\frac{k+5}{2},\frac{k+7}{2}}\Big]$$

for k odd.

A carefull consideration of the topological effect of the braid on the lines L_j shows that they are arranged as in figure 6. To reconstruct the configuration it is enough to know the admissible line l (at angle ϵ w.r.t. the positive real axis). In fact, u_1 is the first point in Π_L (in clockwise order) for k even, or the last in Π_R for k odd. The lines come out of the points in centrifugal directions. They go to infinity, without intersections (so preserving their lexicographical order w.r.t l) with the original asymptotic direction $\frac{\pi}{2} - \epsilon$.

Proposition $1'$ *The matrix representing β' is*

$$A^{\beta'} = \begin{pmatrix}
1 &&&&&&&&&&& \\
& 1 &&&&&&&&&& \\
&& -\binom{k}{k-3} & 1 &&&&&&&& \\
&& -\binom{k}{k-4} & 0 & -\binom{k}{k-5} & 1 &&&&&& \\
&& -\binom{k}{k-5} & 0 & -\binom{k}{k-6} &&&&&&& \\
&& \vdots && \vdots && \cdots &&&&& \\
&& -* && -* && \cdots & 1 &&&& \\
&& -* && -* && \cdots && -\binom{k}{3} & 1 && \\
&& -\binom{k}{\frac{k}{2}-1} && -\binom{k}{\frac{k}{2}-2} && \cdots && -\binom{k}{2} & 0 & -\binom{k}{1} & 1 \\
&& \binom{k}{\frac{k}{2}-2} && \binom{k}{\frac{k}{2}-3} && \cdots && \binom{k}{1} & 0 & 1 & \\
&& * && * && \cdots && 1 &&& \\
&& \vdots && \vdots && \cdots &&&&& \\
&& \binom{k}{2} && \binom{k}{1} &&&&&&& \\
&& \binom{k}{1} & 0 & 1 &&&&&&& \\
&& 1 & 0 &&&&&&&&
\end{pmatrix}$$



Figure 6: Lines L_j (Branch cuts) after the braid which brings S^{-1} to canonoical form

for k even, and

$$A^{\beta'}=\begin{pmatrix}
0 & 1 & & & & & & & & & & & \\
1 & \binom{k}{1} & & & & & & & & & & & \\
 & & 1 & & & & & & & & & & \\
 & & & -\binom{k}{k-4} & 1 & & & & & & & & \\
 & & & -\binom{k}{k-5} & 0 & -\binom{k}{k-6} & 1 & & & & & & \\
 & & & -\binom{k}{k-6} & 0 & -\binom{k}{k-7} & & & & & & & \\
 & & & \vdots & & \vdots & & \cdots & & & & & \\
 & & & -* & & -* & & \cdots & 1 & & & & \\
 & & & -* & & -* & & \cdots & & -\binom{k}{3} & 1 & & \\
 & & & -\binom{k}{\frac{k-3}{2}} & & -\binom{k}{\frac{k-5}{2}} & & \cdots & & -\binom{k}{2} & 0 & -\binom{k}{1} & 1 \\
 & & & \binom{k}{\frac{k-5}{2}} & & \binom{k}{\frac{k-7}{2}} & & \cdots & & \binom{k}{1} & 0 & 1 & \\
 & & & * & & * & & \cdots & & 1 & & & \\
 & & & \vdots & & \vdots & & \cdots & & & & & \\
 & & & \binom{k}{2} & & \binom{k}{1} & & & & & & & \\
 & & & \binom{k}{1} & 0 & 1 & & & & & & & \\
 & & & 1 & 0 & & & & & & & &
\end{pmatrix}$$

for k odd.

The proposition is proved as in the previous section. Finally, the analogous of Theorem 2 holds:

Theorem $2'$ *Consider the Stokes matrix* $S^{-1}=T^{\frac{k}{2}}\ T_F^{-\frac{k}{2}}$ *(k even) or* $S=T^{\frac{k-1}{2}}K_{k2}^{-1}T_F^{\frac{k-1}{2}}$ *(k odd) for the quantum cohomology of* $\mathbf{CP}^{k-1}$ *and set it in the upper triangular form* $S_{upper}^{-1}=P\ S^{-1}\ P^{-1}$ *by the permutation* P*. Then, there exists a braid* β' *(Lemma* $8'$*), represented by a matrix* A^{β} *(Proposition* $1'$*), which sets* S_{upper}^{-1} *in the "canonical form". The entries* s_{ij} *have minus sign for* $j>\frac{k}{2}+1,\ \ i\leq\frac{k}{2}+1$ *and k even, or* $j>\frac{k+3}{2},\ \ i\leq\frac{k+3}{2}$ *and k odd. A suitable conjugation by* $\mathcal{I}=diag(1,1,...1,-1,...,-1)$ *sets all signs positive and the final matrix has entries*

$$s_{ij}=\binom{k}{j-i},\quad s_{ji}=0,\quad \text{for } i<j,\quad s_{ii}=1$$

Proof: The proof is similar to the one of theorem 2.

Examples are found in Appendix 2.

8 Relation between Irregular and Fuchsian systems

Let us consider the fuchsian system

$$(U-\lambda)\ \frac{d\phi}{d\lambda}=\left(\frac{1}{2}+V\right)\ \phi \tag{30}$$

which can also be written

$$\frac{d\phi}{d\lambda}=\sum_{j=1}^{k}\ \frac{A_j}{\lambda-u_j}\ \phi$$

$$A_j=-E_j\ \left(\frac{1}{2}+V\right),\qquad (E_j)_{jj}=1,\quad \text{otherwise } (E_j)_{nm}=0$$

Around the point u_j a fundamental matrix has the form

$$\left[B_0+O\left(\lambda-u_j\right)\right]\ (\lambda-u_j)^{M}$$

where $M=\text{diag}(-\frac{1}{2},0,...,0)$ and the columns of B_0 are the eigenvectors of A_j; in particular, the first column is $(0,..,0,1,0,...,0)^T$, and 1 occurs at the j^{th} position. Then, the system has k independent vector solutions, of which $k-1$ are regular near u_j and the last is

$$\phi^{(j)}(\lambda)=\frac{1}{\sqrt{\lambda-u_j}}\ \begin{pmatrix}0\\ \vdots\\ 1\\ \vdots\\ 0\end{pmatrix}+O\left(\sqrt{\lambda-u_j}\right)\quad \lambda\to u_j$$

where 1 occurs at the j^{th} row. For any u_j we can construct such a basis of solutions. The branch of $\sqrt{\lambda-u_j}$ is chosen as follows: let us consider an angle η with a range of 2π, for example $-\frac{\pi}{2}\leq\eta<\frac{3\pi}{2}$, such that $\eta\neq\arg(u_i-u_j)$, $\forall i\neq j$. Then consider the cuts $L_j=\{\lambda=u_j+\rho e^{i\eta},\ \ \rho>0\}$. Actually, the cuts have two sides, $L_j^{+}=\{\lambda=u_j+\rho e^{i\eta},\ \ \rho>0\}$ and $L_j^{-}=\{\lambda=u_j+\rho e^{i(\eta-2\pi)},\ \ \rho>0\}$. The branch is determined by the choice $\log(\lambda-u_j)=\log|\lambda-u_j|+i\eta$ on L_j^{+} and $\log(\lambda-u_j)=\log|\lambda-u_j|+i(\eta-2\pi)$ on L_j^{-}. On $\mathbf{C}\backslash\bigcup_j L_j$, $\sqrt{\lambda-u_1}$, ..., $\sqrt{\lambda-u_k}$ are single valued.

For any two (column) vector solutions $\phi(\lambda)$, $\psi(\lambda)$ we define the symmetric bilinear form:

$$(\phi,\psi):=\phi(\lambda)^T(\lambda-U)\psi(\lambda)$$

which is independent of λ and u_1, ...,u_k. Let G be the matrix whose entries are $G_{ij}=\left(\phi^{(i)},\phi^{(j)}\right)$. In particular, $G_{ii}=1$. Then, it can be proved (see [2] and also [10]) that near u_j

$$\phi^{(i)}(\lambda)=G_{ij}\phi^{(j)}(\lambda)+r_{ij}(\lambda)$$

where $r_{ij}(\lambda)$ is regular near u_j. For a counter-clockwise loop around u_j the monodromy of $\phi^{(i)}$ is

$$\phi^{(i)}\mapsto R_j\phi^{(i)}:=\phi^{(i)}-2\frac{\left(\phi^{(i)},\phi^{(j)}\right)}{\left(\phi^{(j)},\phi^{(j)}\right)}\ \phi^{(j)}\equiv\phi^{(i)}-2G_{ij}\ \phi^{(j)}$$

Then, the monodromy group of (30) acts on $\phi^{(1)}$, ..., $\phi^{(k)}$ as a reflection group whose Gram matrix is $2G$. In particular, $\phi^{(1)}$, ..., $\phi^{(k)}$ are linearly independent (and then a basis) if and only if $\det G\neq 0$.

Now consider an oriented line l of argument $\theta=\frac{\pi}{2}-\eta$, and for any j define the following vector

$$\tilde{Y}^{(j)}=-\frac{\sqrt{z}}{2\sqrt{\pi}}\int_{\gamma_j}d\lambda\ \phi^{(j)}(\lambda)\ e^{\lambda z} \tag{31}$$

which is a Laplace transform of $\phi^{(j)}$. The path γ_j comes from infinity near L_j^+, encircles u_j and returns to infinity along L_j^-. We can define $\Pi_L = \{\theta < \arg z < \theta + \pi\}$ and $\Pi_R = \{\theta - \pi < \arg z < \theta\}$. $\lambda = \infty$ is a regular singularity for (30), then the integrals exist for $z \in \Pi_L$, and the non-singular matrix $\tilde{Y}(z) := \left[\tilde{Y}^{(1)}|...|\tilde{Y}^{(k)}\right]$ has the asymptotic behaviour

$$\tilde{Y}(z) \sim \left(I + O\left(\frac{1}{z}\right)\right)\ e^{zU} \quad z \to \infty, \quad z \in \Pi_L$$

and satisfies the system (14). Then it is a fundamental matrix $\tilde{Y}_L$. Note that l is admissible, since it does not contain Stokes‘ rays.

It is a fundamental result [10] that the Stokes' matrix of (14) satisfies

$$S + S^T = 2G$$

9 Monodromy Group of the Quantum Cohomology of $\mathbf{CP}^{k-1}$

A system like (30) comes about in the theory of Frobenius manifolds (replace $U \mapsto U(t)$, $V \mapsto V(t)$). It determines flat coordinates $x^1(t,\lambda)$, ..., $x^k(t,\lambda)$ for a linear pencil of metrics $(\ ,\ ) - \lambda <\ ,\ >^*$ ($(\ ,\ )$ is the *intersection form* [9] [10]). We write a gauge equivalent form (gauge $X(t^2)$) at the semisimple point $(0, t^2, 0, ..., 0)$

$$(\hat{\mathcal{U}}(t^2) - \lambda)\ \frac{d\psi}{d\lambda} = \left(\frac{1}{2} + \hat{\mu}\right)\psi \tag{32}$$

A fundamental matrix $\psi(t,\lambda)$ has entries $\psi_\alpha^{(j)}(t,\lambda) = \partial_\alpha x^{(j)}(t,\lambda)$. Moreover, by (31)

$$\partial_\alpha \tilde{t}^j(t,z) = -\frac{\sqrt{z}}{2\sqrt{\pi}} \int d\lambda\ \partial_\alpha x^j(t,\lambda)\ e^{\lambda z} \tag{33}$$

The *Monodromy group* of the Frobenius manifold M is the group of the transformations which $(x^1(t,\lambda), ..., x^k(t,\lambda))$ undergo when t moves in $M \backslash \Sigma_\lambda$, where $\Sigma_\lambda = \{t \in M \mid \det[(\ ,\ ) - \lambda <\ ,\ >] = 0$ $\}$ is the *discriminant* of the linear pencil.

Due to formula (33), for $\mathbf{CP}^{k-1}$ this group is generated by the monodromy of the solutions of (32) when λ describes loops around $u_1(t), ..., u_k(t)$ (see [9] [10]). To these loops, we must add the effect of the displacement $t^2 \mapsto t^2 + 2\pi i$. In fact, in this case

$$[\varphi^{(1)}(ze^{\frac{t^2}{k}}), ..., \varphi^{(k)}(ze^{\frac{t^2}{k}})] \mapsto [\varphi^{(1)}(ze^{\frac{t^2}{k}}), ..., \varphi^{(k)}(ze^{\frac{t^2}{k}})]\ T$$

and the same holds for $\tilde{t}(z, (0, t^2, ..., 0))$.

Then, the monodromy group of the quantum cohomology of $\mathbf{CP}^{k-1}$ is generated by the transformations R_1, R_2, ..., R_k, T introduced in the preceeding sections.

We are going to study the structure of the monodromy group of $\mathbf{CP}^{k-1}$ for any $k \geq 3$. Recall that the matrix S for (14) is not upper triangular, because in U the order of u_1, ..., u_k is not lexicographical w.r.t. the line l. Then, Coxeter identity is $-S^{-1}S^T =$ product of the R_j's in the order refered to l. For example, for $k = 3$, $S^{-1}S^T = -R_2R_3R_1$, since the lexicographical ordering would be u_2, u_3, u_1. From the identity $S^{-1}S^T = (-1)^{k-1}T^k$ it follows a first general relation in the group

$$T^k = (-1)^k \text{ product of } R_j\text{'s in suitable order}$$

Two cases must now be distinguished.

k odd: As a general result [2], $\det G = 0$ if and only if $V + \frac{1}{2}$ has an integer eigenvalue. The eigenvalues of V are $\frac{k-1}{2}$, $\frac{k-3}{2}$, ..., $-\frac{k-1}{2}$. Then, for k odd, $\det G \neq 0$, and $\phi^{(1)}$, ..., $\phi^{(k)}$ are a basis. The matrices R_j are

$$R_j = \begin{pmatrix} 1 & & & & & \\ & \ddots & & & & \\ & & \ddots & & & \\ -2G_{j1} & -2G_{j2} & ... & -1 & ... & -2G_{jk} \\ & & & & \ddots & \\ & & & & & 1 \end{pmatrix}$$

where $S^T + S = 2G$.

In concrete examples, we have "empirically" found other relations like

$$R_2 = p_1(T, R_1)$$

$$R_2 = p_2(T, R_1)$$

$$\vdots$$

$$R_k = p_k(T, R_1)$$

where $p_j(T, R_1)$ means a product of the elements T and R_1. We have also found the relation

$$(TR_1)^k = -I$$

We investigated the following cases:

$\mathbf{CP}^2$ $(k = 3)$

$$R_2 = TR_1T^{-1}, \quad R_3 = T\ (R_1R_2R_1)\ T^{-1}$$
$$(TR_1)^3 = -I$$
$$T^3 = -R_2R_3R_1$$

$\mathbf{CP}^4$ $(k = 5)$

$$\begin{cases} R_2 = TR_1T^{-1}, \quad R_3 = TR_2T^{-1} \\ R_4 = T\ (R_2R_3R_2)\ T^{-1}, \quad R_5 = T^{-1}\ (R_2R_1R_2)\ T \end{cases}$$
$$(TR_1)^5 = -I$$
$$T^5 = -R_3R_4R_2R_5R_1$$

$\mathbf{CP}^6$ $(k = 7)$

$$\begin{cases} R_2 = TR_1T^{-1}, \quad R_3 = TR_2T^{-1}, \quad R_4 = TR_3T^{-1} \\ R_5 = T\ (R_3R_4R_3)T^{-1}, \quad R_6 = T(TR_1)^3\ R_2\ [T(TR_1)^3]^{-1} \\ R_7 = T^{-2}(R_3R_2R_3)T^2 \end{cases}$$
$$(TR_1)^7 = -I$$
$$T^7 = -R_4R_5R_3R_6R_2R_7R_1$$

Note that one ralation, for example that for R_k, can be derived from the others, and that just R_1, T, $-I$ are enought to generate the monodromy group in each of the examples. They satisfy (*in the examples*) the relations:

$$R_1^2 = (-I\ T\ R_1)^k = (-I)^2 = I$$

$$R_1(-I)\ \left((-I)R_1\right)^{-1} = I, \quad T(-I)\ \left((-I)T\right)^{-1} = I$$

The last two relations mean simply the commutativity of $-I$ with R_1 and T. The relations are not only satisfied, but also "fulfilled" (namely, $(-I\ T\ R_1)^n \neq I$ for $n < k$). Now call

$$X := R_1, \quad Y := -ITR_1, \quad Z = -I$$

These elements generate the monodromy group of $\mathbf{CP}^{k-1}$ with *at least* the relations

$$X^2 = Y^k = Z^2 = 1$$

$$(ZX)(XZ)^{-1} = 1, \quad (ZY)(YZ)^{-1} = 1$$

Note that Z generates the cyclic group C_2 of order 2.

If there were no other relations (which we did not find "empirically"), we would conclude that the monodromy group of the quantum cohomology of $\mathbf{CP}^k$ (*in the examples*) is isomorphic to the direct product

$$< X, Y \mid X^2 = Y^k = 1 > \times C_2$$

where $< X, Y \mid X^2 = Y^k = 1 >$ means the group generated by X, Y with relations $X^2 = Y^k = 1$.

k **even:** Now $\det G = 0$, since $V + \frac{1}{2}$ has integer eigenvalues. G has rank $k-1$ and the eigenspace of its eigenvalue 0 has dimension 1. Let $(z^1, ..., z^k)^T$ be an eigenvector of eigenvalue 1. The vector $v := \sum_{j=1}^k z^j \phi^{(j)}$ is zero, because

$$\left(v, \phi^{(i)}\right) = \sum_{j=1}^k z^j G_{ji} = 0 \quad \forall i$$

then

$$z^1 \phi^{(1)} + z^{(2)} \phi^{(2)} + ... + z^k \phi^{(k)} = 0$$

and $k-1$ of the $\phi^{(j)}$'s are linearly independent. The fuchsian system (30) has a regular (vector) solution $\phi_0(\lambda) = \sum_{n=0}^d \phi_n \lambda^n$, where ϕ_n are constant (column) vectors, and ϕ_d is the eigenvector of $V + \frac{1}{2}$ relative to the largest integer eigenvalue less or equal to zero; this eigenvalue is precisely $-d$ (see [2]). In our case, $d = 0$ and $\phi_o(\lambda) = \phi_0$, a constant vector. $\phi^{(1)}$, $\phi^{(2)}$, ..., $\phi^{(k-1)}$, ϕ_0 is then a possible choice for a basis of solutions.

Observe that in the gauge equivalent form $\psi = X\phi$, ψ_0 is the eigenvector of $\frac{1}{2} + \hat{\mu}$ with eigenvalue zero. Then

$$\psi_0 = \begin{pmatrix} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{pmatrix} \equiv \begin{pmatrix} \partial_1 x \\ \partial_2 x \\ \vdots \\ \vdots \\ \partial_k x \end{pmatrix}$$

where all the entries are zero but the one at position $\frac{k}{2} + 1$. x is the flat coordinate for $(\ , \ ) - \lambda < \ , \ >$ corresponding to ψ_0. Then, we can chose the following flat coordinates:

$$x^1(\lambda, t), \ x^2(\lambda, t), \ ... \ , \ x^{k-1}(\lambda, t), \ t^{\frac{k}{2}+1}$$

The monodromy group then acts on a $k-1$ dimensional space.

Let us determine the reduction of R_1, R_2, ..., R_k, T to the $k-1$ dimensional space. The entries of T on the vectors $\phi^{(j)}$ are: $T\phi^{(i)} = \sum_{j=1}^k T_{ji} \phi^{(j)}$, $i = 1, ..., k$. On the new basis $\phi^{(1)}$, ..., $\phi^{(k-1)}$, ϕ_0 the matrices are rewritten

$$R_j \phi^{(i)} = \phi^{(i)} - 2G_{ij} \phi^{(j)} \quad i = 1, .., k-1 \quad j \neq k$$

$$R_j \phi_0 = \phi_0 \qquad j \neq k$$

$$R_k \phi^{(i)} = \phi^{(i)} - 2G_{ik} \left(-\frac{1}{z^k} \sum_{j=1}^{k-1} z^j \phi^{(j)} \right) \quad i \neq k$$

$$T\phi^{(i)} = \sum_{j=1}^{k-1} T_{ji} \phi^{(j)} + T_{ki} \left(-\frac{1}{z^k} \sum_{j=1}^{k-1} z^j \phi^{(j)} \right)$$

Then the matrices assume a reduced form $\begin{pmatrix} * & * & 0 \\ * & * & 0 \\ 0 & 0 & 1 \end{pmatrix}$

We studied two examples; besides Coxeter identity T^k =product of R_j's, we found relations similar to the case k odd:

$$R_2 = p_1(T, R_1)$$

$$R_2 = p_2(T, R_1)$$

$$\vdots$$

$$R_k = p_k(T, R_1)$$

and

$$(TR_1)^k = I$$

Namely:

$\mathbf{CP}^3$ $(k=4)$

$$\begin{cases} R_2 = TR_1T^{-1}, \quad R_3 = TR_2T^{-1} \\ R_4 = T^{-1}\,(R_2R_1R_2)\,T \end{cases}$$

$$(TR_1)^4 = I$$

$$T^4 = R_3R_2R_4R_1$$

$\mathbf{CP}^5$ $(k=6)$

$$\begin{cases} R_2 = TR_1T^{-1}, \quad R_3 = TR_2T^{-1} \\ R_4 = TR_3T^{-1}, \quad R_5 = T\,(R_2R_3R_4R_3R_2)\,T^{-1} \\ R_6 = T^{-1}\,(R_2R_1R_2)\,T \end{cases}$$

$$(TR_1)^6 = I$$

$$T^6 = R_4R_3R_5R_2R_6R_1$$

The same remarks of k odd hold here. Call

$$X := R_1, \quad Y := R_1T$$

then, if there were no other hiden relations, the monodromy group of the quantum cohomology of $\mathbf{CP}^k$ (*in the examples*) would be isomorphic to

$$< X, Y, \mid X^2 = Y^k = 1 >$$

Note that $< X, Y, \mid X^2 = Y^k = 1 >$ is (isomorphic to) the subgroup of orientation preserving transformations of the hyperbolic triangular group $[2, k, \infty]$.

Lemma 10: *The subgroup of the orientation preserving transformations of the hyperbolic triangular group* $[2, k, \infty]$ *is isomorphic to the subgroup of* $PSL(2, \mathbf{R})$ *generated by*

$$\tau \mapsto -\frac{1}{\tau}$$

$$\tau \mapsto \frac{1}{2\cos\frac{\pi}{k} - \tau}$$

$\tau \in H := \{z \in \mathbf{C} \mid \Im z > 0\}$

Proof: Consider three integers m_1, m_2, m_3 such that

$$\frac{1}{m_1} + \frac{1}{m_2} + \frac{1}{m_3} < 1$$

In the Bolyai-Lobatchewsky plane H, the triangular group $[m_1, m_3, m_3]$ of hyperpolic reflections in the sides of hyperbolic triangles of angles $\frac{\pi}{m_1}$, $\frac{\pi}{m_2}$, $\frac{\pi}{m_3}$ is generated by three reflections r_1, r_2, r_3 satisfying the relations

$$r_1^2 = r_2^2 = r_3^2 = (r_2r_3)^{m_1} = (r_3r_1)^{m_2} = (r_2r_1)^{m_3} = 1$$

and the subgroup of orientation preserving transformation is generated by $X = r_2r_3$, $Y = r_3r_1$, Then

$$X^{m_1} = Y^{m_2} = (XY)^{m_3} = 1$$

For $m_1 = 2$, $m_2 = k$, $m_3 = \infty$, a fundamental triangular region is $\{0 < \Re z < \cos\frac{\pi}{k}\} \cap \{|z| > 1\}$. Then

$$r_1(\tau) = -\bar{\tau}, \quad r_2(\tau) = \frac{1}{\bar{\tau}}, \quad r_3(\tau) = 2\cos\frac{\pi}{k} - \bar{\tau}$$

The bar means complex conjugation. Then

$$X(\tau) = -\frac{1}{\tau}, \quad Y(\tau) = \frac{1}{2\cos\frac{\pi}{k} - \tau}$$

$\square$

Remark : The orientation preserving transformations of $[2, 3, \infty]$ are the *modular group* $PSL(2, \mathbf{Z})$.

Theorem 3: *The monodromy group of the quantum cohomology of* $\mathbf{CP}^2$ *is isomorphic to*

$$< X, Y, \ | \ X^2 = Y^3 = 1 > \times C_2 \cong PSL(2, \mathbf{Z}) \times C_2 \tag{34}$$

The monodromy group of the quantum cohomology of $\mathbf{CP}^3$ *is isomorphic to*

$$< X, Y, \ | \ X^2 = Y^4 = 1 > \cong \ \textit{orient. preserv. transf. of } [2, 4, \infty] \tag{35}$$

The theorem for the case of $\mathbf{CP}^2$ is already proved in [10].

Proof : a) $\mathbf{CP}^2$:

$$R_1 = \begin{pmatrix} -1 & 3 & 3 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \qquad T = \begin{pmatrix} 0 & 0 & 1 \\ -1 & 3 & 3 \\ 0 & -1 & 0 \end{pmatrix}$$

and $X = R_1$, $Y = -ITR_1$ and $Z = -I$ satisfy the relations of (34). They act on the column vector $\mathbf{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix}$. The quadratic form $q(x, y, z) = \mathbf{x}^T G \mathbf{x}$ is R_1 and T - invariant. Then T, R_1 act on two dimensional invariant subspaces $q(x, y, z) = \text{constant}$. On each of these subspaces we introduce new coordinates $\chi \in \mathbf{R}$ and $\varphi \in [0, 2\pi)$. Let $\tau = e^{\chi} \ e^{i\varphi}$ and

$$x = \frac{a}{2}(\tau\bar{\tau} - \frac{3}{2}(\tau + \bar{\tau}) + 1) \ \frac{i}{\tau - \bar{\tau}}$$

$$y = \frac{a}{2}(\tau\bar{\tau} - \frac{1}{2}(\tau + \bar{\tau}) - 1) \ \frac{i}{\tau - \bar{\tau}}$$

$$z = \frac{a}{2}(-\tau\bar{\tau} - \frac{1}{2}(\tau + \bar{\tau}) + 1) \ \frac{i}{\tau - \bar{\tau}}$$

$a \in \mathbf{R}$, $a \neq 0$. Note that $q(x, y, z) = a^2 > 0$. Then, it is easily verified that

$$\mathbf{x}(-\frac{1}{\tau}) = -X \ \mathbf{x}(\tau)$$

$$\mathbf{x}(\frac{1}{1 - \tau}) = Y \ \mathbf{x}(\tau)$$

$$\mathbf{x}(\tau, -a) = Z \ \mathbf{x}(\tau, a)$$

. This implies the 1 to 1 correspondence between the generators of the modular group and X and Y.

b) Case of $\mathbf{CP}^3$.

$$R_1 = \begin{pmatrix} -1 & 4 & -10 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad T = \begin{pmatrix} 0 & 0 & 1 & 0 \\ -1 & 0 & 3 & 0 \\ 0 & -1 & 3 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

The matrices are already written on $\phi^{(1)}$, $\phi^{(2)}$, $\phi^{(3)}$, ϕ_0. Recall that the monodromy acts only on x^1, x^2, x^3, because the last flat coordinate is t^3. This action is given by the following three dimensional matrices, acting on a three dimensional space of vectors $\mathbf{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix}$

$$r_1 = \begin{pmatrix} -1 & 4 & 10 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad t := \begin{pmatrix} 0 & 0 & 1 \\ -1 & 0 & 3 \\ 0 & -1 & 3 \end{pmatrix}$$

We redefine $X = r_1$ and $Y = tr_1$, which satisfy the relations (35). We proceed as above, defining

$$x = a(\tau\bar{\tau} - \frac{1}{\sqrt{2}}(\tau + \bar{\tau}) + \frac{1}{3}) \ \frac{i}{\tau - \bar{\tau}}$$

$$y = a(-\frac{2}{3}\tau\bar{\tau} - \frac{2\sqrt{2}}{3}(\tau + \bar{\tau}) + \frac{2}{3})\ \frac{i}{\tau - \bar{\tau}}$$

$$z = a(-\frac{1}{3}\tau\bar{\tau} - \frac{\sqrt{2}}{6}(\tau + \bar{\tau}) + \frac{1}{3})\ \frac{i}{\tau - \bar{\tau}}$$

$a \neq 0$. Note that $\mathbf{x}^T g \mathbf{x} = (8/9)a^2$, where g is the 3×3 reduction of G. It is easily verified that

$$\mathbf{x}(-\frac{1}{\tau}) = -X\ \mathbf{x}(\tau)$$

$$\mathbf{x}(\frac{1}{\sqrt{2} - \tau}) = -Y\ \mathbf{x}(\tau)$$

which proves the theorem.

□

APPENDIX 1: Proof of Lemma 5

Let us consider the function

$$g(z) = C \int_{-c-i\infty}^{-c+i\infty} ds\ \Gamma^k(-s)\ e^{i\pi f s}\ z^{ks}$$

where C and f are constants to be determined later, $c > 0$. The path of integration is a vertical line through $-c$.

a) Domain of definition. We use Stirling formula

$$\Gamma(-s) = e^s e^{-(s+\frac{1}{2})\log(-s)}\sqrt{2\pi}(1 + O(1/s)) \qquad s \to \infty, \quad |\arg(-s)| < \pi$$

where $\log(-s) = \log(s) + i\pi$.

The integrand is

$$\Gamma^k(-s)e^{i\pi f s}z^{ks} \sim e^{ks(1+\log z)}e^{-k(s+\frac{1}{2})\log s}e^{-ik\pi(s+\frac{1}{2})+i\pi s f} \qquad s \to \infty$$

Now let $s = -c + i\eta$, $ds = i d\eta$. The integral in η is on the real axis. The dominant part in the integrand is

$$e^{\eta[k\arg(i\eta - c) + k\pi - k\arg z - \pi f]}$$

The condition of (uniform) convergence of the integral is

$$-\frac{\pi}{2} - f\frac{\pi}{k} < \arg z < \frac{\pi}{2} - f\frac{\pi}{k}$$

b) g solves (13). In fact (for simplicity $C = 1$ here):

$$(z\partial_z)^k g(z) = k^k \int_{-c-i\infty}^{-c+i\infty} ds\ \Gamma^k(-s)e^{i\pi f s}s^k z^{ks} = (-1)^k k^k \int_{-c-i\infty}^{-c+i\infty} ds\ \Gamma^k(1-s)e^{i\pi f s}z^{ks}$$

where we have used the identity $s\Gamma(-s) = -\Gamma(1-s)$. Now let $t = s - 1$:

$$(z\partial_z)^k g(z) = (-1)^k e^{i\pi f}(zk)^k \int_{-c-1-i\infty}^{-c-1+i\infty} dt\ \Gamma^k(-t)e^{i\pi f t}z^{kt}$$

$$= (-1)^k e^{i\pi f}(kz)^k g(z)$$

Now we impose $(-1)^k e^{i\pi f} = 1$, namely

$$f = k + 2m \qquad m \in \mathbf{Z}$$

Point b) is proved.

c) Asymptotic behaviour of $g(z)$. We use Laplace method for analytic functions (the so called "steepest descent method"). Put $C = 1$. By Stirling's formula

$$g(z) = (2\pi)^{k/2} e^{-i\frac{\pi}{2}k} \int_{-c-i\infty}^{-c+i\infty} ds\ e^{\phi(s)}$$

$$\phi(s) = s\ [k(1+\log z) + i\pi f - i\pi k] - k\left(s+\frac{1}{2}\right)\log s + O\left(\frac{1}{s}\right)$$

The the stationary point of $\phi(s)$ is

$$s_0 = z e^{-i\frac{\pi}{k}(k-f)} - \frac{1}{2} + O\left(\frac{1}{z}\right) \qquad z \to \infty$$

In the hypothesis $z \to \infty$ a straightforward computation gives

$$\phi(s_0) \sim kze^{-i\frac{\pi}{k}(k-f)} - \frac{k}{2}\log z + i\frac{\pi}{2}(k-f) \qquad \frac{d^2\phi(s_0)}{ds^2} \sim -\frac{k}{z}e^{i\frac{\pi}{k}(k-f)}$$

We are ready to apply the steepest descent method. We deform the path of integration in such a way that it passes through s_0. Let us call it γ.

$$g(z) = (2\pi)^{\frac{k}{2}} e^{-i\frac{\pi}{2}k} e^{\phi(s_0)} \int_\gamma ds\ e^{\phi(s)-\phi(s_0)}$$

$$\sim (2\pi)^{\frac{k}{2}} e^{-i\frac{\pi}{2}k} e^{\phi(s_0)} \int_\gamma ds\ e^{\frac{1}{2}\frac{d^2\phi(s_0)}{ds^2}(s-s_0)^2}$$

Let us divide γ in two paths: γ_1 from s_0 to $+i\infty$ and γ_2 from $-i\infty$ to s_0. The integrals becomes the sum of two integrals. In $\int_{\gamma_1}$ we change variable. Let $\tau > 0$ and

$$-\tau^2 = \frac{1}{2}\frac{d^2\phi(s_0)}{ds^2}(s-s_0)^2 = -\frac{k}{2z}\left[e^{i\frac{\pi}{k}(k-f)}(s-s_0)^2\right]$$

Then

$$\int_{\gamma_1} e^{\frac{d^2\phi(s_0)}{ds^2}(s-s_0)^2} = \frac{1}{2}\sqrt{\frac{\pi}{k}}(2z)^{\frac{1}{2}} e^{-i\frac{\pi}{2k}(k-f)-i\pi} + O(e^{-\alpha|z|}) \qquad \alpha > 0$$

Note that $\int_{\gamma_1} = \int_{\gamma_2}$. Now, recalling that $f = k + 2m$, we conclude that

$$g(z) \sim (2\pi)^{\frac{k+1}{2}} e^{-i(m+1)\pi} e^{-i\frac{\pi}{2}k}\ \frac{1}{\sqrt{k}}\frac{e^{i\frac{\pi}{k}m}}{z^{\frac{k-1}{2}}}\ \exp(ke^{i\frac{2\pi}{k}m}z) \qquad z \to \infty$$

If we choose

$$m = n-1+kl \qquad l \in \mathbf{Z}$$

and $C = \left[(2\pi)^{\frac{k+1}{2}} e^{i\pi(l-\frac{k}{2}-n-kl)}\right]^{-1}$ we have

$$g(z) \sim \frac{1}{\sqrt{k}}\frac{e^{i\frac{\pi}{k}(n-1)}}{z^{\frac{k-1}{2}}}\exp(ke^{i\frac{2\pi}{k}(n-1)}z)$$

Then g is a solution $\varphi^{(n)}$ on the domain

$$-\frac{3\pi}{2} - 2(n-1)\frac{\pi}{k} - 2\pi l < \arg z < -\frac{\pi}{2} - 2(n-1)\frac{\pi}{k} - 2\pi l$$

d) We now determine the domain where the analytic extension of $g(z)$ still has the above asymptotic behaviour. From Lemma 4 we derived that the first entries of the $n(k)^{th}$ columns of Y_L and Y_R are equal and coincide with $\varphi_L^{(n(k))}(z) \equiv \varphi_R^{(n(k))}(z)$ times $z^{\frac{k-1}{2}}$, and $\varphi^{(n(k))}$ has the estabilished asymptotic behaviour on the enlarged domains

$$-\frac{\pi}{k} - \pi < \arg z < \pi + \frac{\pi}{k}$$

$$-\pi < \arg z < \pi + \frac{2\pi}{k}$$

If in $g(z)$ we chose $n = n(k)$ and $l = -1$ the integral representation holds for $-\frac{\pi}{2} < \arg z < \frac{\pi}{2}$ for k even, and for $-\frac{\pi}{2} + \frac{\pi}{k} < \arg z < \frac{\pi}{2} + \frac{\pi}{k}$ for k odd. Then its analytic continuation is precisely $\varphi_L^{(n(k))}(z) \equiv \varphi_R^{(n(k))}(z)$.

e) We prove the identity (16). Observe that $(z\partial_z)^k \varphi = (kz)^k \varphi$ is invariant for $z \mapsto z\, e^{\frac{2\pi i}{k}}$. A generic solution can be reperesented near $z = 0$ as

$$\varphi(z) = \sum_{m=0}^{\infty} \frac{z^{km}}{(m!)^k} \left[a_m^{(1)} + a_m^{(2)} \log z + ... + a_m^{(k)} \log^{k-1} z\right]$$

It follows that the operator $(A\varphi)(z) = \varphi(ze^{\frac{2\pi i}{k}})$ has eigenvalues 1, because on the basis obtained with $(a_0^{(1)} = 1,\ a_0^{(2)} = ... = a_0^{(k)} = 0)$, $(a_0^{(1)} = 0,\ a_0^{(2)} = 1,\ a_0^{(3)} = ... = a_0^{(k)} = 0)$, ..., $(a_0^{(1)} = ... = a_0^{(k-1)} = 0,\ a_0^{(k)} = 1)$ it is represented by a lower triangular matrix having 1's on the diagonal. Then $(A-1)^k = 0$ and

$$(A-1)^k g(z) = 0$$

is precisely our identity. Lemma 4 is proved.

□

APPENDIX 2:

First we give Φ_R and Φ_L. k **odd:**

$$\Phi_R(z)^T = \begin{bmatrix} (-1)^{\frac{k-1}{2}} \left[g(ze^{-\frac{2\pi i}{k}(\frac{k-1}{2})}) - \binom{k}{1} g(ze^{-\frac{2\pi i}{k}(\frac{k-3}{2})}) + \ldots + \binom{k}{k-1} g(ze^{\frac{2\pi i}{k}(\frac{k-1}{2})})\right] \\ \vdots \\ g(ze^{-\frac{4\pi i}{k}}) - \binom{k}{1} g(ze^{-\frac{2\pi i}{k}}) + \binom{k}{2} g(z) - \binom{k}{3} g(ze^{\frac{2\pi i}{k}}) + \binom{k}{4} g(ze^{\frac{4\pi i}{k}}) \\ -g(ze^{-\frac{2\pi i}{k}}) + \binom{k}{1} g(z) - \binom{k}{2} g(ze^{\frac{2\pi i}{k}}) \\ g(z) \\ -g(ze^{\frac{2\pi i}{k}}) \\ g(ze^{\frac{4\pi i}{k}}) \\ \vdots \\ (-1)^{\frac{k-1}{2}} g(ze^{\frac{2\pi i}{k}(\frac{k-1}{2})}) \end{bmatrix}$$

$$\Phi_L(z)^T = \begin{bmatrix} (-1)^{\frac{k-1}{2}} g(ze^{-\frac{2\pi i}{k}(\frac{k-1}{2})}) \\ -(-1)^{\frac{k-1}{2}} g(ze^{-\frac{2\pi i}{k}(\frac{k-3}{2})}) \\ \vdots \\ -g(ze^{-\frac{2\pi i}{k}}) \\ g(z) \\ -g(ze^{\frac{2\pi i}{k}}) + \binom{k}{k-1} g(z) \\ g(ze^{\frac{4\pi i}{k}}) - \binom{k}{k-1} g(ze^{\frac{2\pi i}{k}}) + \binom{k}{k-2} g(z) - \binom{k}{k-3} g(ze^{-\frac{2\pi i}{k}}) \\ \vdots \\ (-1)^{\frac{k-1}{2}} \left[g(ze^{\frac{2\pi i}{k}(\frac{k-1}{2})}) - \binom{k}{k-1} g(ze^{\frac{2\pi i}{k}(\frac{k-3}{2})}) + \ldots - \binom{k}{2} g(ze^{-\frac{2\pi i}{k}(\frac{k-3}{2})})\right] \end{bmatrix}$$

k **even:**

$$\Phi_R(z)^T = \begin{bmatrix} (-1)^{\frac{k}{2}}\left[g(ze^{-i\pi}) - \binom{k}{1} g(ze^{-i\pi+i\frac{2\pi}{k}}) + \ldots + \binom{k}{k-1} g(ze^{i(\pi-\frac{2\pi}{k})})\right] \\ \vdots \\ g(ze^{-\frac{4\pi i}{k}}) - \binom{k}{1} g(ze^{-\frac{2\pi i}{k}}) + \binom{k}{2} g(z) - \binom{k}{3} g(ze^{\frac{2\pi i}{k}}) \\ -g(ze^{-\frac{2\pi i}{k}}) + \binom{k}{1} g(z) \\ g(z) \\ -g(ze^{\frac{2\pi i}{k}}) \\ g(ze^{\frac{4\pi i}{k}}) \\ \vdots \\ (-1)^{\frac{k}{2}-1} g(ze^{\frac{2\pi i}{k}(\frac{k}{2}-1)}) \end{bmatrix}$$

$$\Phi_L(z)^T = \begin{bmatrix} (-1)^{\frac{k}{2}} g(ze^{-i\pi}) \\ \vdots \\ g(ze^{-i\frac{4\pi}{k}}) \\ -g(ze^{-\frac{2\pi i}{k}}) \\ g(z) \\ -g(ze^{\frac{2\pi i}{k}}) + \binom{k}{k-1} g(z) - \binom{k}{k-2} g(ze^{-i\frac{2\pi}{k}}) \\ g(ze^{\frac{4\pi i}{k}}) - \binom{k}{k-1} g(ze^{\frac{2\pi i}{k}}) + \binom{k}{k-2} g(z) - \binom{k}{k-3} g(ze^{-\frac{2\pi i}{k}}) + \binom{k}{k-4} g(ze^{-i\frac{4\pi}{k}}) \\ \vdots \\ (-1)^{\frac{k}{2}-1}\left[g(ze^{i\pi-i\frac{2\pi}{k}}) - \binom{k}{k-1} g(ze^{i\pi-i\frac{4\pi}{k}}) + \ldots - \binom{k}{2} g(ze^{-i\pi+i\frac{2\pi i}{k}})\right] \end{bmatrix}$$

We give all the matrices of interest up to $k = 10$. S_{upper} is PSP^{-1}. A stends for A^{β}, A' for $A^{\beta'}$. $S^{\beta} = AS_{upper}A^T$, $S^{\beta'} = A'S^{-1}_{upper}[A']^T$. $\sigma_{i,i+1} := \beta^{-1}_{i,i+1}$

CP^2

$$K_{12} = \begin{pmatrix} 1 & 0 & 0 \\ -3 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad K_{13} := \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -3 & 0 & 1 \end{pmatrix} \quad K_{32} := \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 3 \\ 0 & 0 & 1 \end{bmatrix}$$

$$T := \begin{bmatrix} 0 & 0 & 1 \\ -1 & 3 & 3 \\ 0 & -1 & 0 \end{bmatrix}$$

$$S := \begin{bmatrix} 1 & 0 & 0 \\ -3 & 1 & 3 \\ -3 & 0 & 1 \end{bmatrix} \quad PSP^{-1} = \begin{bmatrix} 1 & 3 & -3 \\ 0 & 1 & -3 \\ 0 & 0 & 1 \end{bmatrix}$$

CP^3

$$K_{42} := \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 6 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad K_{43} := \begin{bmatrix} 1 & 0 & 0 & 0 \\ -4 & 1 & 0 & 0 \\ 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$T := \begin{bmatrix} 0 & 0 & 0 & 1 \\ -1 & 0 & 6 & 4 \\ 0 & -1 & 4 & 0 \\ 0 & 0 & -1 & 0 \end{bmatrix} \quad T_1 := \begin{bmatrix} 0 & 0 & 1 & 0 \\ -1 & 0 & 3 & 0 \\ 0 & -1 & 3 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$S := \begin{bmatrix} 1 & 0 & 0 & 0 \\ -4 & 1 & 0 & 6 \\ 10 & -4 & 1 & -20 \\ -4 & 0 & 0 & 1 \end{bmatrix}$$

$$P := \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{bmatrix} \quad S_{upper} = \begin{bmatrix} 1 & -4 & -20 & 10 \\ 0 & 1 & 6 & -4 \\ 0 & 0 & 1 & -4 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$A := \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 4 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad A' = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -4 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix} \quad S^{\beta} = S^{\beta'} := \begin{bmatrix} 1 & 4 & 6 & -4 \\ 0 & 1 & 4 & -6 \\ 0 & 0 & 1 & -4 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

where $\beta = \beta_{12}$

$\mathbf{CP}^4$

$$K_{52} := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ -5 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 10 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \quad K_{53} := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 10 \\ 0 & 0 & 1 & 5 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$T := \begin{bmatrix} 0 & 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 10 & 5 \\ 0 & -1 & 5 & 10 & 0 \\ 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \end{bmatrix} \quad S = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ -5 & 1 & 0 & 0 & 10 \\ 15 & -5 & 1 & 5 & -40 \\ 40 & -10 & 0 & 1 & -95 \\ -5 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$P := \begin{bmatrix} 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 \end{bmatrix} \quad S_{upper} := \begin{bmatrix} 1 & 5 & -5 & -40 & 15 \\ 0 & 1 & -10 & -95 & 40 \\ 0 & 0 & 1 & 10 & -5 \\ 0 & 0 & 0 & 1 & -5 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$A := \begin{bmatrix} 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 5 & 0 & 0 \\ 0 & 1 & 10 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \quad A' := \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 1 & 5 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -5 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

$$S^{\beta} = S^{\beta'} := \begin{bmatrix} 1 & 5 & 10 & 10 & -5 \\ 0 & 1 & 5 & 10 & -10 \\ 0 & 0 & 1 & 5 & -10 \\ 0 & 0 & 0 & 1 & -5 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

where $\beta = \beta_{23}\beta_{12}$, $\beta' = \beta_{12}\sigma_{45}$.

$\mathbf{CP}^5$

$$K_{62} := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 15 \\ 0 & 0 & 1 & 0 & 15 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \quad K_{63} := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ -6 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 20 \\ 0 & 0 & 0 & 1 & 6 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$T := \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 & 15 & 6 \\ 0 & -1 & 0 & 15 & 20 & 0 \\ 0 & 0 & -1 & 6 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 \end{bmatrix} \quad S := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ -6 & 1 & 0 & 0 & 0 & 15 \\ 21 & -6 & 1 & 0 & 15 & -70 \\ -56 & 21 & -6 & 1 & -84 & 210 \\ 105 & -20 & 0 & 0 & 1 & -294 \\ -6 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$T_1 := \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 0 \\ -1 & 0 & 0 & 0 & 5 & 0 \\ 0 & -1 & 0 & 15 & 25 & 0 \\ 0 & 0 & -1 & 6 & 1 & 0 \\ 0 & 0 & 0 & -1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$P := \begin{bmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$A := \begin{bmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 6 & 0 & 0 \\ 1 & 6 & 0 & 15 & 0 & 0 \\ 0 & 0 & 1 & 20 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \quad A' := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -20 & 1 & 0 & 0 \\ 0 & 0 & -15 & 0 & -6 & 1 \\ 0 & 0 & 6 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{bmatrix}$$

$$S^{\beta} := \begin{bmatrix} 1 & 6 & 15 & 20 & 15 & -6 \\ 0 & 1 & 6 & 15 & 20 & -15 \\ 0 & 0 & 1 & 6 & 15 & -20 \\ 0 & 0 & 0 & 1 & 6 & -15 \\ 0 & 0 & 0 & 0 & 1 & -6 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \quad S^{\beta'} := \begin{bmatrix} 1 & 6 & 15 & 20 & -15 & -6 \\ 0 & 1 & 6 & 15 & -20 & -15 \\ 0 & 0 & 1 & 6 & -15 & -20 \\ 0 & 0 & 0 & 1 & -6 & -15 \\ 0 & 0 & 0 & 0 & 1 & 6 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

where $\beta = \beta_{12}(\beta_{34}\beta_{23}\beta_{12})$, $\beta' = [(\sigma_{34}\sigma_{56})\sigma_{45}]\sigma_{56}$.

$\mathbf{CP}^6$

$$K_{72} := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 21 \\ 0 & 0 & 1 & 0 & 0 & 35 & 0 \\ 0 & 0 & 0 & 1 & 7 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \quad K_{73} := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -7 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 35 \\ 0 & 0 & 0 & 1 & 0 & 21 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$T := \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 & 0 & 21 & 7 \\ 0 & -1 & 0 & 0 & 35 & 35 & 0 \\ 0 & 0 & -1 & 7 & 21 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 \end{bmatrix} \quad S := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -7 & 1 & 0 & 0 & 0 & 0 & 21 \\ 28 & -7 & 1 & 0 & 0 & 35 & -112 \\ -84 & 28 & -7 & 1 & 7 & -224 & 378 \\ -378 & 112 & -21 & 0 & 1 & -728 & 1638 \\ 224 & -35 & 0 & 0 & 0 & 1 & -728 \\ -7 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$P := \begin{bmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$A := \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 7 & 0 & 0 \\ 1 & 0 & 7 & 0 & 21 & 0 & 0 \\ 0 & 1 & 21 & 0 & 35 & 0 & 0 \\ 0 & 0 & 0 & 1 & 35 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \quad A' := \begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 7 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -35 & 1 & 0 & 0 \\ 0 & 0 & 0 & -21 & 0 & -7 & 1 \\ 0 & 0 & 0 & 7 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{bmatrix}$$

$$S^{\beta} := \begin{bmatrix} 1 & 7 & 21 & 35 & 35 & 21 & -7 \\ 0 & 1 & 7 & 21 & 35 & 35 & -21 \\ 0 & 0 & 1 & 7 & 21 & 35 & -35 \\ 0 & 0 & 0 & 1 & 7 & 21 & -35 \\ 0 & 0 & 0 & 0 & 1 & 7 & -21 \\ 0 & 0 & 0 & 0 & 0 & 1 & -7 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \quad S^{\beta'} := \begin{bmatrix} 1 & 7 & 21 & 35 & 35 & -21 & -7 \\ 0 & 1 & 7 & 21 & 35 & -35 & -21 \\ 0 & 0 & 1 & 7 & 21 & -35 & -35 \\ 0 & 0 & 0 & 1 & 7 & -21 & -35 \\ 0 & 0 & 0 & 0 & 1 & -7 & -21 \\ 0 & 0 & 0 & 0 & 0 & 1 & 7 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

where $\beta = (\beta_{23}\beta_{12})(\beta_{45}\beta_{34}\beta_{23}\beta_{12})$, $\beta' = \beta_{12}[(\sigma_{45}\sigma_{67})\sigma_{56}]\sigma_{67}$.

$\mathbf{CP}^7$

$$K_{82} := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 28 \\ 0 & 0 & 1 & 0 & 0 & 0 & 70 & 0 \\ 0 & 0 & 0 & 1 & 0 & 28 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \quad K_{83} := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -8 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 56 \\ 0 & 0 & 0 & 1 & 0 & 0 & 56 & 0 \\ 0 & 0 & 0 & 0 & 1 & 8 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$T := \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 & 0 & 0 & 28 & 8 \\ 0 & -1 & 0 & 0 & 0 & 70 & 56 & 0 \\ 0 & 0 & -1 & 0 & 28 & 56 & 0 & 0 \\ 0 & 0 & 0 & -1 & 8 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \end{bmatrix}$$

$$S := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -8 & 1 & 0 & 0 & 0 & 0 & 0 & 28 \\ 36 & -8 & 1 & 0 & 0 & 0 & 70 & -168 \\ -120 & 36 & -8 & 1 & 0 & 28 & -504 & 630 \\ 330 & -120 & 36 & -8 & 1 & -216 & 2100 & -1848 \\ -1512 & 378 & -56 & 0 & 0 & 1 & -3912 & 7476 \\ 420 & -56 & 0 & 0 & 0 & 0 & 1 & -1560 \\ -8 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$P := \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \qquad S_{upper} := \begin{bmatrix} 1 & -8 & -216 & 36 & 2100 & -120 & -1848 & 330 \\ 0 & 1 & 28 & -8 & -504 & 36 & 630 & -120 \\ 0 & 0 & 1 & -56 & -3912 & 378 & 7476 & -1512 \\ 0 & 0 & 0 & 1 & 70 & -8 & -168 & 36 \\ 0 & 0 & 0 & 0 & 1 & -56 & -1560 & 420 \\ 0 & 0 & 0 & 0 & 0 & 1 & 28 & -8 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & -8 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$A := \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 8 & 0 & 0 \\ 0 & 1 & 0 & 8 & 0 & 28 & 0 & 0 \\ 1 & 8 & 0 & 28 & 0 & 56 & 0 & 0 \\ 0 & 0 & 1 & 56 & 0 & 70 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 56 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \qquad A' := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -56 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -70 & 0 & -56 & 1 & 0 & 0 \\ 0 & 0 & -56 & 0 & -28 & 0 & -8 & 1 \\ 0 & 0 & 28 & 0 & 8 & 0 & 1 & 0 \\ 0 & 0 & 8 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$S^{\beta} := \begin{bmatrix} 1 & 8 & 28 & 56 & 70 & 56 & 28 & -8 \\ 0 & 1 & 8 & 28 & 56 & 70 & 56 & -28 \\ 0 & 0 & 1 & 8 & 28 & 56 & 70 & -56 \\ 0 & 0 & 0 & 1 & 8 & 28 & 56 & -70 \\ 0 & 0 & 0 & 0 & 1 & 8 & 28 & -56 \\ 0 & 0 & 0 & 0 & 0 & 1 & 8 & -28 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & -8 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \qquad S^{\beta'} := \begin{bmatrix} 1 & 8 & 28 & 56 & 70 & -56 & -28 & -8 \\ 0 & 1 & 8 & 28 & 56 & -70 & -56 & -28 \\ 0 & 0 & 1 & 8 & 28 & -56 & -70 & -56 \\ 0 & 0 & 0 & 1 & 8 & -28 & -56 & -70 \\ 0 & 0 & 0 & 0 & 1 & -8 & -28 & -56 \\ 0 & 0 & 0 & 0 & 0 & 1 & 8 & 28 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 8 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

where $\beta = (\beta_{34}\beta_{23}\beta_{12})\beta_{23}(\beta_{56}\beta_{45}\beta_{34}\beta_{23}\beta_{12})$, $\beta' = [(\sigma_{34}\sigma_{56}\sigma_{78})(\sigma_{45}\sigma_{67})\sigma_{56}][(\sigma_{67}\sigma_{78})\sigma_{67}]$.

$\mathbf{CP}^8$

$$K_{92} := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 36 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 126 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 84 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 9 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \qquad K_{93} := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -9 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 84 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 126 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 36 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$T := \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 36 & 9 \\ 0 & -1 & 0 & 0 & 0 & 0 & 126 & 84 & 0 \\ 0 & 0 & -1 & 0 & 0 & 84 & 126 & 0 & 0 \\ 0 & 0 & 0 & -1 & 9 & 36 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \end{bmatrix}$$

$$S := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -9 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 36 \\ 45 & -9 & 1 & 0 & 0 & 0 & 0 & 126 & -240 \\ -165 & 45 & -9 & 1 & 0 & 0 & 84 & -1008 & 990 \\ 495 & -165 & 45 & -9 & 1 & 9 & -720 & 4620 & -3168 \\ 3168 & -990 & 240 & -36 & 0 & 1 & -3015 & 25740 & -19932 \\ -4620 & 1008 & -126 & 0 & 0 & 0 & 1 & -15867 & 25740 \\ 720 & -84 & 0 & 0 & 0 & 0 & 0 & 1 & -3015 \\ -9 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$P := \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$S_{upper} := \begin{bmatrix} 1 & 9 & -9 & -720 & 45 & 4620 & -165 & -3168 & 495 \\ 0 & 1 & -36 & -3015 & 240 & 25740 & -990 & -19932 & 3168 \\ 0 & 0 & 1 & 84 & -9 & -1008 & 45 & 990 & -165 \\ 0 & 0 & 0 & 1 & -126 & -15867 & 1008 & 25740 & -4620 \\ 0 & 0 & 0 & 0 & 1 & 126 & -9 & -240 & 45 \\ 0 & 0 & 0 & 0 & 0 & 1 & -84 & -3015 & 720 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 36 & -9 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & -9 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$A := \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 9 & 0 & 0 \\ 0 & 0 & 1 & 0 & 9 & 0 & 36 & 0 & 0 \\ 1 & 0 & 9 & 0 & 36 & 0 & 84 & 0 & 0 \\ 0 & 1 & 36 & 0 & 84 & 0 & 126 & 0 & 0 \\ 0 & 0 & 0 & 1 & 126 & 0 & 126 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 84 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \qquad A' := \begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 9 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -126 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -126 & 0 & -84 & 1 & 0 & 0 \\ 0 & 0 & 0 & -84 & 0 & -36 & 0 & -9 & 1 \\ 0 & 0 & 0 & 36 & 0 & 9 & 0 & 1 & 0 \\ 0 & 0 & 0 & 9 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$S^{\beta} = \begin{bmatrix} 1 & 9 & 36 & 84 & 126 & 126 & 84 & 36 & -9 \\ 0 & 1 & 9 & 36 & 84 & 126 & 126 & 84 & -36 \\ 0 & 0 & 1 & 9 & 36 & 84 & 126 & 126 & -84 \\ 0 & 0 & 0 & 1 & 9 & 36 & 84 & 126 & -126 \\ 0 & 0 & 0 & 0 & 1 & 9 & 36 & 84 & -126 \\ 0 & 0 & 0 & 0 & 0 & 1 & 9 & 36 & -84 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 9 & -36 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & -9 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \qquad S^{\beta'} := \begin{bmatrix} 1 & 9 & 36 & 84 & 126 & 126 & -84 & -36 & -9 \\ 0 & 1 & 9 & 36 & 84 & 126 & -126 & -84 & -36 \\ 0 & 0 & 1 & 9 & 36 & 84 & -126 & -126 & -84 \\ 0 & 0 & 0 & 1 & 9 & 36 & -84 & -126 & -126 \\ 0 & 0 & 0 & 0 & 1 & 9 & -36 & -84 & -126 \\ 0 & 0 & 0 & 0 & 0 & 1 & -9 & -36 & -84 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 9 & 36 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 9 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

where $\beta = (\beta_{45}\beta_{34}\beta_{23}\beta_{12})(\beta_{34}\beta_{23})(\beta_{67}\beta_{56}\beta_{45}\beta_{34}\beta_{23}\beta_{12})$, and $\beta' = \beta_{12}[(\sigma_{45}\sigma_{67}\sigma_{89})(\sigma_{56}\sigma_{78})\sigma_{67}][(\sigma_{78}\sigma_{89})\sigma_{78}]$.

$\mathbf{CP}^9$

$$K_{10,2} := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 45 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 210 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 210 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 45 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$K_{10,3} := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -10 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 120 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 252 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 120 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 10 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$T := \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 45 & 10 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 210 & 120 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 210 & 252 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 45 & 120 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 10 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \end{bmatrix}$$

$$S := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -10 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 45 \\ 55 & -10 & 1 & 0 & 0 & 0 & 0 & 0 & 210 & -330 \\ -220 & 55 & -10 & 1 & 0 & 0 & 0 & 210 & -1848 & 1485 \\ 715 & -220 & 55 & -10 & 1 & 0 & 45 & -1980 & 9240 & -5148 \\ -2002 & 715 & -220 & 55 & -10 & 1 & -440 & 10395 & -34320 & 15015 \\ 17160 & -4752 & 990 & -120 & 0 & 0 & 1 & -25190 & 177705 & -120120 \\ -11880 & 2310 & -252 & 0 & 0 & 0 & 0 & 1 & -52910 & 73755 \\ 1155 & -120 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & -5390 \\ -10 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$P := \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$S_{upper} := \begin{bmatrix} 1 & -10 & -440 & 55 & 10395 & -220 & -34320 & 715 & 15015 & -2002 \\ 0 & 1 & 45 & -10 & -1980 & 55 & 9240 & -220 & -5148 & 715 \\ 0 & 0 & 1 & -120 & -25190 & 990 & 177705 & -4752 & -120120 & 17160 \\ 0 & 0 & 0 & 1 & 210 & -10 & -1848 & 55 & 1485 & -220 \\ 0 & 0 & 0 & 0 & 1 & -252 & -52910 & 2310 & 73755 & -11880 \\ 0 & 0 & 0 & 0 & 0 & 1 & 210 & -10 & -330 & 55 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & -120 & -5390 & 1155 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 45 & -10 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & -10 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$A = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 10 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 10 & 0 & 45 & 0 & 0 \\ 0 & 1 & 0 & 10 & 0 & 45 & 0 & 120 & 0 & 0 \\ 1 & 10 & 0 & 45 & 0 & 120 & 0 & 210 & 0 & 0 \\ 0 & 0 & 1 & 120 & 0 & 210 & 0 & 252 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 252 & 0 & 210 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 120 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$A' := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -120 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -210 & 0 & -252 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -252 & 0 & -210 & 0 & -120 & 1 & 0 & 0 \\ 0 & 0 & -210 & 0 & -120 & 0 & -45 & 0 & -10 & 1 \\ 0 & 0 & 120 & 0 & 45 & 0 & 10 & 0 & 1 & 0 \\ 0 & 0 & 45 & 0 & 10 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 10 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$S^{\beta} := \begin{bmatrix} 1 & 10 & 45 & 120 & 210 & 252 & 210 & 120 & 45 & -10 \\ 0 & 1 & 10 & 45 & 120 & 210 & 252 & 210 & 120 & -45 \\ 0 & 0 & 1 & 10 & 45 & 120 & 210 & 252 & 210 & -120 \\ 0 & 0 & 0 & 1 & 10 & 45 & 120 & 210 & 252 & -210 \\ 0 & 0 & 0 & 0 & 1 & 10 & 45 & 120 & 210 & -252 \\ 0 & 0 & 0 & 0 & 0 & 1 & 10 & 45 & 120 & -210 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 10 & 45 & -120 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 10 & -45 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & -10 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$S^{\beta'} := \begin{bmatrix} 1 & 10 & 45 & 120 & 210 & 252 & -210 & -120 & -45 & -10 \\ 0 & 1 & 10 & 45 & 120 & 210 & -252 & -210 & -120 & -45 \\ 0 & 0 & 1 & 10 & 45 & 120 & -210 & -252 & -210 & -120 \\ 0 & 0 & 0 & 1 & 10 & 45 & -120 & -210 & -252 & -210 \\ 0 & 0 & 0 & 0 & 1 & 10 & -45 & -120 & -210 & -252 \\ 0 & 0 & 0 & 0 & 0 & 1 & -10 & -45 & -120 & -210 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 10 & 45 & 120 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 10 & 45 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 10 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

where $\beta = (\beta_{56}\beta_{45}\beta_{34}\beta_{23}\beta_{12})(\beta_{45}\beta_{34}\beta_{23})\beta_{34}(\beta_{78}\beta_{67}\beta_{56}\beta_{45}\beta_{34}\beta_{23}\beta_{12})$,
and $\beta' = [(\sigma_{34}\sigma_{56}\sigma_{78}\sigma_{9,10})(\sigma_{45}\sigma_{67}\sigma_{89})(\sigma_{56}\sigma_{78})\sigma_{67}][(\sigma_{78}\sigma_{89}\sigma_{9,10})(\sigma_{78}\sigma_{89})\sigma_{78}]$.

References

[1] W.Balser, W.B.Jurkat, D.A.Lutz: *Birkhoff Invariants and Stokes' Multipliers for Meromorphic Linear Differential Equations*, Journal Math. Analysis and Applications, **71**, (1979), 48-94.

[2] W.Balser, W.B.Jurkat, D.A.Lutz: *On the reduction of connection problems for differential equations with an irregular singular point to ones with only regular singularities*, SIAM J. Math. Anal., **12**, (1981), 691-721.

[3] Beilinson A.A. : *Coherent Sheaves on* $\mathbf{P}^n$ *and Problems of Linear Algebra*, Functional Anal. Appl., **12**, (1978).

[4] J.S.Birman: *Braids, Links, and Mapping Class Groups*, Annals of Mathematics Studies, **82**, Princeton Univ. Press (1975).

[5] Bondal A.I., Polishchuk A.E. : *Homologival Properties of Associative Algebras: the Method of Helices*, Russ.Acad.Sci.Izv.Math., **42**, (1994), 219-260

[6] Cecotti S. Vafa C.: *On Classification of $N = 2$ supersymmetric theories*,Comm.Math.Phys, **158**, (1993), 569-644.

[7] B.Dubrovin: *Integrable Systems in Topological Field Theory*, Nucl. Phys B, **379**, (1992), 627-689.

[8] B.Dubrovin: *Geometry and Itegrability of Topological-Antitopological Fusion*, Comm.Math.Phys, **152**, (1993), 539-564.

[9] B.Dubrovin: *Geometry of 2D topological field theories*, Lecture Notes in Math, **1620**, (1996), 120-348.

[10] B.Dubrovin: *Painlevé trascendents in two-dimensional topological field theory*, SISSA Preprint 24/98/FM, (1998).

[11] B.Dubrovin: *Geometry and Analytic Theory of Frobenius Manifolds*, math.AG/9807034, (1998).

[12] B.Dubrovin- M.Mazzocco: *Monodromy of Certain Painlevé trascendents and Reflection Groups*, SISSA Preprint 149/97/FM - To appear in ..., (1997).

[13] M.Jimbo, T.Miwa, K.Ueno : *Monodromy Preserving Deformations of Linear Ordinary Differential Equations with Rational Coefficients (I)*, Physica , **D 2** , (1981), 306

[14] M.Jimbo, T.Miwa : *Monodromy Preserving Deformations of Linear Ordinary Differential Equations with Rational Coefficients (II)*, Physica , **D 2** , (1981), 407

[15] M.Jimbo, T.Miwa : *Monodromy Preserving Deformations of Linear Ordinary Differential Equations with Rational Coefficients (III)*, Physica , **D 4** , (1981), 26

[16] Gorodentsev A.L., Rudakov A.N. : Duke Mathematical Journal, **54**, (1987), 115

[17] A.R.Its - V.Y.Novokshenov: *The isomonodromic deformation method in the theory of Painleve equations*, Lecture Notes in Math, **1191**, (1986).

[18] Kontsevich M.: Talk at the Scuola Normale Superiore. Pisa, April 1998

[19] Kontsevich M., Manin Y.I. : *Gromow-Witten classes, Quantum Cohomology and Enumerative Geometry*, Comm.Math.Phys, **164**, (1994), 525-562

[20] Manin V.I. : *Frobenius Manifolds, Quantum Cohomology and Moduli Spaces*, Max Planck Institut fur Mathematik. Bonn. Germany, (1998 ?).

[21] McDuff D., Salamon D. : *J-holomorphic Curves and Quantum Cohomology*, Providence RI. American Mathematical Society, (1994)

[22] Ruan Y., Tian G. : *A Mathematical Theory of Quantum Cohomology*, Math.Res.Lett., **1**, (1994), 269-278

[23] Rudakov A.: *Integer Valued Bilinear Forms and Vector Bundles*, Math. USSR Sbornik, **66**, (1990), 187-194.

[24] Rudakov A. : *Helices and Vector Bundles*, Seminaire Rudakov. London Math. Society. Lecture notes Series, **148**, C.U.P (1990).

[25] Zaslow E.: *Solitons and Helices: The Search for a Math-Phys Bridge* , Comm.Math.Phys, **175**, (1996), 337-375.